



PENERBIT  **PRESS**
UNIVERSITI TEKNOLOGI MARA

eISSN: 2785-924X
2023 March, Volume 3, Issue 1

JOURNAL OF **SMART SCIENCE AND TECHNOLOGY**



Vol. 3 No. 1 (2023): Journal of Smart Science and Technology



[Issue Information](#)

***Note: Please take note that the DOIs will take some time to be deposited after publication. Your patience and understanding are highly appreciated. Thank you.*

DOI: <https://doi.org/10.24191/jsst.v3i1>

Published: 2023-03-30

Articles

Simulation of a Pumped Stormwater System and Evaluation of the Solar Potential for Pumping

Hasan Huseyin Coban

1-13

 [Download](#)

The Effect of the Kaolinitic Clay and Asphaltenes on the Rheological Properties of Trinidad Lake Asphalt and Trinidad Petroleum Bitumen-Clay Composites

Rean Maharaj

14-24

 download

Exploring Bipartite Network Approach in Hand, Foot and Mouth Disease Hotspot Identification

Chin Ying Liew, Nor Shamira Sabri, Boon Hao Hong, Jane Labadin

25-36

 download

Effects of Cellulolytic Fungi and Biofertilizer Doses on the Growth and Yield of Tomato (*Solanum lycopersicum* L.) in Alluvial Soil

Agus Suyanto, Agnes Tutik Purwani Irianti, Setiawan

37-43

 download

Analysis of Development Planning of Own Use Gas Pipeline from Gas Compressor Stations to Gathering Stations in X Field

Muhammad Ariyon, Muhammad Setriya Ramadhan

44-53

 download

Clinical Efficacy of Autogenous Vertebrae-Filled PEEK Cage in Anterior Cervical Discectomy and Fusion with Instrumentation

Ahmad Nazrun Shuid, Mohd Fairudz Mohd Miswan, Xuehua Zhan, Wenqi Yuan, Haifeng Yuan

54-62

 download

Performance of Correlational Filtering and Deep Learning Based Single Target Tracking Algorithms

ZhongMing Liao, Azlan Ismail

63-79

 download

Platform &
workflow by
OJS / PKP

[Home](#) / Editorial Team

Editorial Team

Chief Editor: Professor Ts. Dr. Khong Heng Yen (Faculty of Applied Sciences, Universiti Teknologi MARA, Malaysia) (khonghy@uitm.edu.my)

Managing Editor: Associate Professor Dr. Robin Chang Yee Hui (Faculty of Applied Sciences, Universiti Teknologi MARA, Malaysia) (robincyh@uitm.edu.my)

Co-Managing Editor: Associate Professor Dr. Leong Siow Hoo (Faculty of Computer and Mathematical Sciences, Universiti Teknologi MARA, Malaysia) (leongsh@uitm.edu.my)

Copy Editor: Dr. Liew Chin Ying (Faculty of Computer and Mathematical Sciences, Universiti Teknologi MARA, Malaysia) (cyliew@uitm.edu.my)

Advisory Board:

- Professor Dato Dr. Jamil Hamali (Universiti Teknologi MARA, Malaysia) jamil1@uitm.edu.my
- Professor Dr. Haeng-Ki Lee (KAIST, South Korea) haengki@kaist.ac.kr
- Professor Dr. Yoshinori Asakawa (Tokushima Bunri University, Japan) asakawa@ph.bunri-u.ac.jp

Editors:

- Professor Ir. Ts. Dr. Mohamad Rusop Mahmood (Center for Functional Materials & Nanotechnology, Institute of Science, Universiti Teknologi MARA, Malaysia) rusop@uitm.edu.my
- Professor Ir. Ts. Dr. Yupiter Harangan Prasada Manurung (Smart Manufacturing Research Institute, Universiti Teknologi MARA, Malaysia) yupiter.manurung@uitm.edu.my
- Professor Dr. Yap Bee Wah (Universiti Teknologi MARA, Malaysia) yapbeewah@uitm.edu.my
- Professor Dr. Yana M Syah (Faculty of Mathematics and Natural Sciences, Institut Teknologi Bandung, Indonesia) yana@chem.itb.ac.id
- Professor Dr. Zhang Lei (Anhui Medical University, China) zhangl@ahmu.edu.cn
- Professor Dr. Harikesh Bahadur Singh (GLA University, India) hb.singh@gla.ac.in
- Assistant Professor Dr. Wongi S NA (Seoul National University of Science & Technology, South Korea) wongi@seoultech.ac.kr
- Assistant Professor Dr. Hammad R. Khalid, King Fahd University of Petroleum & Minerals, Saudi Arabia) hammad.khalid@kfupm.edu.sa
- Associate Professor Dr. Siva K Balasundram (Faculty of Agriculture, Universiti Putra Malaysia, Malaysia) siva@upm.edu.my

- Associate Professor Dr. Bo Xia (Queensland University of Technology, Australia) paul.xia@qut.edu.au
- Associate Professor Dr. Huang Zhiyong (School of Computing, National University of Singapore, Singapore) dcshuang@nus.edu.sg
- Associate Professor Ir. Ts. Dr Mohamad Hafiz Mamat (NANO-ElecTronic Centre (NET), School of Electrical Engineering, College of Engineering, Universiti Teknologi MARA, Malaysia) mhmamat@uitm.edu.my
- Associate Professor Dr. Sim Siong Fong (Faculty of Resource Science and Technology, Universiti Sarawak Malaysia, Malaysia) sfsim@unimas.my
- Dr. Jesús del Barrio Lasheras (Instituto de Nanociencia y Materiales de Aragón, Universidad de Zaragoza, Spain) jdb529@unizar.es
- Dr. Rudy Tawie Ak Joseph Sipi (College of Engineering, Universiti Teknologi MARA, Malaysia) rudy@uitm.edu.my
- Dr. Ahmad Faiz Abdul Rashid (Faculty of Architecture, Planning and Surveying, Universiti Teknologi MARA, Malaysia) afaiz@uitm.edu.my
- ChM. Dr. Cindy Tan Soo Yun (Faculty of Applied Sciences, Universiti Teknologi MARA, Malaysia) cindytan@uitm.edu.my

Associate Editors

- Dr. Asmah Alia Mohamad Bohari (Faculty of Architecture, Planning and Surveying, Universiti Teknologi MARA, Malaysia) asmahalia@uitm.edu.my
- Associate Professor Dr. Clotilda Petrus (College of Engineering, Universiti Teknologi MARA, Malaysia) clotilda166@uitm.edu.my
- Dr. Liew Fui Kiew (Faculty of Applied Sciences, Universiti Teknologi MARA, Malaysia) fkiew@uitm.edu.my
- Associate Professor Dr. Wan Fairos Wan Yaacob (Faculty of Business and Management, Universiti Teknologi MARA, Malaysia) wnfairos@uitm.edu.my

Platform &
workflow by
OJS / PKP

Effects of *Cellulolytic* Fungi and Biofertilizer Doses on the Growth and Yield of Tomato (*Solanum lycopersicum* L.) in Alluvial Soil

Agus Suyanto^{1,*}, Agnes Tutik Purwani Irianti¹, Setiawan¹

¹Agrotechnology Study Program, Faculty of Agriculture, Universitas Panca Bhakti, Kom Yos Sudarso Street, Pontianak City 78113, Indonesia

Received: 29-09-2022

Revised: 02-12-2022

Accepted: 13-12-2022

Published: 30-03-2023

*Correspondence

Email: agussuyanto@upb.ac.id

(Agus Suyanto)

DOI: <https://doi.org/10.24191/jsst.v3i1.36>

© 2023 The Author(s). Published by UiTM Press. This is an open access article under the terms of the Creative Commons Attribution 4.0 International Licence (<http://creativecommons.org/licenses/by/4.0/>), which permits use, distribution and reproduction in any medium, provided the original work is properly cited.



Abstract

The production and harvest areas of tomatoes in West Kalimantan tend to decrease. One of the cultivation techniques that is expected to increase the yield and quality of the plant is by using organic fertilizers. The study objective was to determine the effects of interactions between cellulolytic fungi and biofertilizer doses on the growth and yield of tomato (*Solanum lycopersicum* L.) in alluvial soils. A completely randomized design (CRD) with factorial patterns was applied in the research. The first factor which was cellulolytic fungi in biofertilizers (J) included: J0 (organic fertilizer without decomposer), J1 (organic fertilizer with *Aspergillus* sp.), J2 (organic fertilizer with *Trichoderma* spp.), and J3 (organic fertilizer with *Aspergillus* sp. and *Trichoderma* spp.). The second factor that was of biofertilizer doses (D) consisted of: D1 (25 g/polybag), D2 (50 g/polybag), D3 (75 g/polybag), and D4 (100 g/polybag). The results showed no significant difference between the cellulolytic fungi and biofertilizer doses on all observation variables ($p > 0.05$). There was no significant main effect of cellulolytic fungus type on all observation variables, while the treatment of biofertilizer doses had a significant effect on plant height, total leaf number per plant, fruit number per plant, and fruit weight per plant. The J3D4 treatment gave the best results on the plant height (97.67 cm) and the leaf number (18.17 strands), while the J2D4 treatment gave the best results on the fruit number per plant (3.31 fruits), and the fruit weight per plant (13.32 g). The addition of cellulolytic fungi to biofertilizers at a dose of 100g/polybag can increase growth and yield on tomatoes in alluvial soil.

Keywords

Aspergillus sp.; *Trichoderma* sp.; Cellulolytic fungi; Biofertilizer; *Solanum lycopersicum* L.

Citation: Suyanto, A., Irianti, A. T. P., & Setiawan. (2023). Effects of cellulolytic fungi and biofertilizer doses on the growth and yield of tomato (*Solanum lycopersicum* L.) in alluvial soil. *Journal of Smart Science and Technology*, 3(1), 37-43.

1 Introduction

Tomato (*Solanum lycopersicum* L.) is one of the most sought-after horticultural commodities, and it is needed in various

places in Indonesia as tomato is consumed as vegetable and fruit. However, both production and harvest areas of the plant in West Kalimantan tend to decrease. The tomato yields of the province from 2016 to

2018 were 3,766 tons of 426 ha, 1,805 tons of 369 ha, and 1,128 tons of 374 ha¹. The decline was led by the increasing pest pressure and intensified chemical treatments. It was also due to the reduced soil fertility especially its organic matter and phosphorus which are the major constraints that limit crop production. The high pressure on agricultural land also reduced the availability of soil nutrients and caused a significant reduction in soil fertility and crop yields^{2,3}.

To increase tomato productivity, there are many things which can be performed from improving cultivation techniques to postharvest treatment. To improve the cultivation techniques, one of the most important things to focus on is the availability of sufficient nutrients, as nutrition for the plant growth and development will then affect the quality and quantity of yielded tomatoes. One of the cultivation techniques that is expected to increase the yield and quality of the plant is by using organic fertilizers. The use of biofertilizers along with inorganic fertilizers could be the right answer since biofertilizers were eco-friendly and more economic, and it played an important role in reducing the dependence on chemical fertilizers⁴. Rice straw is one of the materials which can be used as organic fertilizer.

Rice straw is a crop waste that is generated in vast amount during harvest time. Rice straw was a part of a rice plant that includes stems, leaves, and tillers⁵. Composting rice straws would provide a method to avoid air pollution that comes from residual burning while also preventing the loss of nutrients in organic materials. The waste is a local material that can potentially be turned into organic fertilizer. Rice straws were abundantly available during harvest, yet they have not been fully utilized⁶. Recycling plant waste by composting is important for maintaining soil fertility in tropical regions. Global production of rice straws is estimated to be around 650-975 million tonnes annually. There is about 1-1.5 kg of rice straws from each kilogram of shifted rice produced. Various studies found that rice straws contain nutrients: N, P, K, and C-organic, which are needed by soil and plants. Rice straw contains around 0.6% N, 1.5% K,

5% Si, 40% C, and 0.10% of each P and S. The utilization of crop waste as compost would not only provide plants with important nutrients but also increase soil fertility, and it can be a method to protect the environment through waste management⁵.

Rice straws are a potential source of soil organic matter (SOM). However, the high C/N ratio contained is the main obstacle if it is directly applied to soil as SOM. In addition to reducing the volume of material to facilitate application, composting rice straw is required to avoid negative effects on plants. Composting process was regulated by the fundamental environmental factors such as temperature, moisture content, pH, aeration and some characteristics such as C/N ratio, particle size and nutrients of organic wastes⁷. The higher the C/N ratio, the longer the rate of decomposition, and vice versa. The availability of decomposers is one of factors which also determines the composting rate, so it is necessary to find effective decomposers to reduce the C/N ratio of crop waste to be used as compost.

Decomposer is a biological activator used to accelerate composting process and to improve compost quality. The number and type of microorganisms also determine the success of the composting process⁸. In ecosystems, decomposing microorganisms play a critical role because the remains of dead organisms will be broken down into elements in the form of nutrients: N, P, K, Ca, and Mg to be returned to soil and/or in the form of gases: CH₄ or CO₂ to be released into the atmosphere. The use of decomposers which are in accordance with organic substrates and soil conditions is an effective alternative used for decomposition acceleration and fertilization supplementation⁹. Organic decomposers consist of a group of fungi and bacteria. Cellulolytic fungi as organic decomposer generally have a better ability than bacteria in decomposing plant remains (hemicellulose, cellulose, and lignin). *Aspergillus* sp. and *Trichoderma* spp. are fungi that can be used as biofertilizers because they are able to decompose and inhibit the growth of fungal plant pathogens.

In accordance to this, it is important to conduct a study to obtain fungal isolates

which have a high ability in degrading organic matter so as to optimize the utilization of rice straws as organic fertilizer which can be biological agents for plant protection. This study aimed to determine the effects of interactions between cellulolytic fungi and biofertilizer doses on the growth and yield of tomato (*Solanum lycopersicum* L.) in alluvial soils.

2 Materials and Method

The research was conducted at the Green House of Faculty of Agriculture, Panca Bhakti University, Pontianak, for a duration of four months, from May 2020 to August 2020. The materials included tomato seeds of Tymoti F1 variety, *Trichoderma* spp., *Aspergillus* sp., alluvial soils, polybags, biofertilizers, rice straw compost and inorganic fertilizers (urea, SP-36, and KCL), and dolomite lime (CaMg (CO₃)₂). The tools used consisted of hoes, machetes, shovels, ordinary scales, analytical scales, thermometers, hygrometers, meters, soil sieves, cameras, stationery, and other supporting tools.

A completely randomized design (CRD) with factorial patterns was applied in the research. The first factor that includes decomposer types in organic fertilizers (J) consisted of four treatment levels, while the second factor that was organic fertilizer doses (D) consisted of four treatment levels. Each treatment was repeated three times and each repetition involved three plants; therefore, there were 144 plants used altogether. The treatment details were as follows.

- a. The first factor: cellulolytic fungi in biofertilizers (J), namely:

J0: organic fertilizer without decomposer
 J1: organic fertilizer with *Aspergillus* sp.
 J2: organic fertilizer with *Trichoderma* spp.
 J3: organic fertilizer with *Aspergillus* sp. and *Trichoderma* spp.

- b. The second factor: biofertilizer doses (D), namely:

D1: 25 g/polybag
 D2: 50 g/polybag
 D3: 75 g/polybag
 D4: 100 g/polybag

The treatment combinations were: J₀D₁, J₀D₂, J₀D₃, J₀D₄, J₁D₁, J₁D₂, J₁D₃, J₁D₄, J₂D₁, J₂D₂, J₂D₃, J₂D₄, J₃D₁, J₃D₂, J₃D₃, and J₃D₄.

2.1 Research Implementation

2.1.1 Preparation of Compost Materials

This process took a month. Rice straws, manure, bran, and other additives were used for the treatment of compost materials.

2.1.2 Preparation of Planting Media

The planting media used were alluvial soil taken from a depth of ± 20 cm. The soils were washed, dried, and sieved before being placed in a 10 kg polybag.

2.1.3 Seed Preparation and Seeding

Polybags with alluvial soil and rice husk charcoal 2 : 1 were used. Alluvial soils used in the study belong to the classification of the order Inceptisol. The physical properties of alluvial soils are steep and shapeless, with slow permeability. The chemical properties of alluvial soils are low organic matter content and low soil pH. Before seeding, rice seeds were selected and immersed in water for 15 minutes; the floating seeds were discarded and the sunk seeds were used and seeded.

2.1.4 Liming

Liming was done three weeks before planting, and it aimed to increase the soil pH to 6.0 using dolomite lime at a dose of 9.6 grams/polybag.

2.1.5 Fertilization

- a. The fertilizers used in this research consisted of biofertilizers, rice straw compost, urea, SP-36, and KCL.
- b. The biofertilizers were made from rice straw compost that had been treated for one month with decomposers *Trichoderma* spp. and *Aspergillus* sp. Fertilizer was applied seven days before planting. The decomposer

types and biofertilizer doses were acceptable for the treatments.

- c. Inorganic fertilizers were administered at the following rates: 0.7 g/polybag urea, 1.25 g/polybag SP-36, and 1 g/polybag KCL. Fertilizer was applied during planting.

2.1.6 Transplanting of Tomato Seedlings into Polybags

The seedlings were transplanted into the polybags after 21-23 days of seeding. Thinning was only carried out when there were unhealthy or dead plants, no later than two weeks after planting.

2.1.7 Staking

To keep the plants upright, bamboo or wooden stakes were installed when the plants were 21 days after planting. The stakes were installed at a distance of 5 cm from the tomato plants with a minimum depth of 10 cm.

2.1.8 Maintenance

- a. Watering

It was carried out twice a day, in the morning and the afternoon, taking into account the moisture of the media.

- b. Weed control

The weeds surrounding the plants in the polybags were pulled out mechanically.

2.1.9 Harvesting

The tomatoes were harvested after 66 days of being transplanted (DAT) into the polybags. They were harvested after ripening with physiological characteristics: the change of fruit skin from light green to pink or to dark red. Tomatoes were harvested as many as six times in the morning.

2.2 Research Observations

The parameters observed were :

- a. Plant height (cm)

It was measured at the end of the research from the base of the trunk to the highest leaf tip.

- b. Leaf number per plant (strand)

The number of leaves was calculated for those that had been completely formed at the end of the research.

- c. Fruit number per plant (unit)

It was calculated by summing the total fruit produced from each treatment that was carried out six times during harvesting.

- d. Fruit weight per plant (gram)

The weight was calculated with scales by calculating the total weight of fruit produced from each treatment that was carried out six times during harvesting.

2.3 Data Analysis

The *F*-test at the 5% level of significance was performed to determine the effects of all treatments. If there was a significant effect on the observed parameters, each treatment would be compared using honestly significant difference (HSD) test at 5%.

3 Results and Discussion

Based on the statistical analysis, the interaction of cellulolytic fungi and biofertilizer doses did not have a significant effect on all observation variables. The treatment of single cellulolytic fungus type also did not show any significant effect on all observation variables, while the treatment of biofertilizer doses had a significant effect on plant height and a highly significant effect on leaf number per plant, fruit number per plant, and fruit weight per plant. The J3D4 treatment gave the best results on the plant height (97.67 ± 3.54 cm) and the leaf number (18.17 ± 1.15 strands), while the J2D4 treatment gave the best results on the fruit number per plant (3.31 ± 0.17 fruits), and

the fruit weight per plant (13.32 ± 1.36 g) (Table 1).

The application of bio-fertilizers which were the compost enriched with *Trichoderma* spp. and *Aspergillus* sp. was suspected to be able to maintain the neutrality of soil pH and to contribute to the increase of the total soil C, N, P, and K, and the tomato yields. The incorporation of biofertilizers in soil plays a major role in improving soil fertility, yield attributing characters and final yield. Biofertilizers

were certainly known to enhance the nutrient availability for crop plants and to impart better health to plants and soils, thus enhancing crop yields as they are modernized forms of organic fertilizers into which beneficial microorganisms have been incorporated¹⁰. The best plant growth was through the application of organic fertilizers, which provide an additional source of nutrients and improve the efficiency of mineral fertilizers by making nutrients more available for plant growth¹¹.

Table 1. Results of the average effects of *Cellulolytic* fungi and biofertilizer doses on growth and yield of tomato (*Solanum lycopersicum* L.) in alluvial soil.

No.	Treatments	Plant height \pm SD (cm)	Leaf number/Plant \pm SD (strand)	Fruit number/Plant* \pm SD (unit)	Fruit weight/Plant* \pm SD (gram)
1	J0D1	78.67 \pm 4.64	13.83 \pm 0.76	2.73 \pm 0.19	11.13 \pm 0.41
2	J0D2	85.06 \pm 3.99	16.39 \pm 0.91	3.04 \pm 0.15	12.37 \pm 0.17
3	J0D3	87.11 \pm 16.51	16.11 \pm 1.57	3.35 \pm 0.53	13.59 \pm 1.27
4	J0D4	95.56 \pm 11.23	18.56 \pm 4.24	3.33 \pm 0.24	14.13 \pm 1.26
5	J1D1	83.33 \pm 5.77	13.00 \pm 1.73	1.32 \pm 1.41	3.94 \pm 5.95
6	J1D2	81.61 \pm 7.05	15.11 \pm 1.01	2.79 \pm 0.32	11.70 \pm 1.60
7	J1D3	91.17 \pm 14.91	16.50 \pm 1.50	3.16 \pm 0.38	11.94 \pm 1.20
8	J1D4	94.00 \pm 12.00	16.06 \pm 1.41	3.04 \pm 0.25	12.79 \pm 1.28
9	J2D1	82.56 \pm 16.22	14.44 \pm 1.26	2.06 \pm 1.35	7.59 \pm 6.14
10	J2D2	85.56 \pm 4.85	16.22 \pm 1.34	2.81 \pm 0.29	12.14 \pm 0.67
11	J2D3	83.00 \pm 4.72	15.22 \pm 2.03	2.75 \pm 0.49	12.48 \pm 0.22
12	J2D4	90.78 \pm 18.37	17.83 \pm 1.04	3.31 \pm 0.17	13.32 \pm 1.36
13	J3D1	84.94 \pm 7.87	15.56 \pm 1.38	2.57 \pm 0.58	9.67 \pm 2.29
14	J3D2	87.22 \pm 3.34	15.00 \pm 1.45	3.20 \pm 0.22	13.02 \pm 1.59
15	J3D3	80.83 \pm 9.25	15.56 \pm 2.34	3.22 \pm 0.18	12.73 \pm 0.69
16	J3D4	97.67 \pm 3.54	18.17 \pm 1.15	2.91 \pm 0.05	12.00 \pm 0.26

Note: * = Transformation Yield Data, $\sqrt{x} + 0.5$; SD = Standard deviation; $n = 48$

Symbiotic relationships between fungi and a variety of plants can occur, and this can produce colonies on the outside part of a root system. This condition could increase the uptake of water and nutrients by the plant roots¹². The result could be linked to the presence of *Trichoderma* spp. and *Aspergillus* sp. in the compost. *Trichoderma* spp. certainly can grow into the soil and exert spatial and nutrient competition. *Trichoderma* spp. had the ability to release bioactive molecules and facilitate the supply of nutrients¹³. Sawadogo et al.¹⁴ tested the effect of compost enriched

with *Trichoderma harzianum* on tomato plants and showed that the high content of total N, P and K in biofertilizers could be due to the fungus that has the capacity to colonize the environment in which it was found to promote the mineralization of organic matter.

Aspergillus sp. is known to be able to absorb phosphorus for its own nutritional requirements, and this promotes the availability of the soluble form of phosphorus for easy absorption by plant roots. In the solubilization of phosphorus for nutrient plants, *Aspergillus* sp. was

identified as an active biological agent to promote the availability of soluble phosphorus in the soil¹⁵. *Aspergillus* sp. was also known to produce plant growth promoting substances such as indole-3-acetic acid (IAA) and gibberellic acid (GA) that affect plant growth parameters and nutrient uptake¹⁶.

4 Conclusion

In accordance with the research results, the interaction of cellulolytic fungi and biofertilizer doses did not have a significant effect on all observation variables. The treatment of single cellulolytic fungus type also did not show any significant effect on all observation variables, while the treatment of biofertilizer doses had a significant effect on plant height and a highly significant effect on leaf number per plant, fruit number per plant, and fruit weight per plant. The J3D4 treatment gave the best results on the plant height (97.67 cm) and the leaf number (18.17 strands), while the J2D4 treatment gave the best results on the fruit number per plant (3.31 fruits), and the fruit weight per plant (13.32 g).

Conflict of Interest

The authors reported that there was no conflict of interest in this work.

Acknowledgment

The authors are grateful to the Agriculture Faculty, Universitas Panca Bhakti, Pontianak for the support and assistance.

Funding

This research was funded by the authors themselves.

Author Contribution

Conceptualization: Suyanto, A.
Data curation: Suyanto, A.
Methodology: Suyanto, A., Irianti, A.T.P., & Setiawan
Formal analysis: Setiawan
Visualisation: Irianti, A.T.P.
Software: Suyanto, A.
Writing (original draft): Suyanto, A., Irianti, A.T.P., & Setiawan

Writing (review and editing): Suyanto, A., Irianti, A.T.P., & Setiawan

Validation: Suyanto, A., Irianti, A.T.P., & Setiawan

Supervision: Suyanto, A.

Funding acquisition: Irianti, A.T.P.

Project administration: Setiawan

References

1. Kementerian Pertanian Republik Indonesia. (2019). *Luas panen dan produksi tomat menurut provinsi tahun 2015-2019*. <https://www.pertanian.go.id>
2. Son, D., Somda, I., Legreve, A., & Schiffrers, B. (2018). Effect of plant diversification on pest abundance and tomato yields in two cropping systems in Burkina Faso: Farmer practices and integrated pest management. *International Journal of Biological and Chemical Sciences*, 12(1), 101-119. <https://doi.org/10.4314/ijbcs.v12i1.8>
3. Coulibaly, P. J. A., Sawadogo, J., Bambara, Y. A. I., Ouédraogo, W. B. M., Legma, J. B., & Compaoré, E. (2021). Effect of bio-fertilizers on tomato (*Solanum lycopersicum*) production and on soil physico-chemical properties in Sudan area of Burkina Faso. *Current Agriculture Research Journal*, 9(1), 43-53. <https://doi.org/10.12944/CARJ.9.1.06>
4. Kamal, S., Kumar, M., Rajkumar, & Raghav, M. (2018). Effect of Biofertilizers on Growth and Yield of Tomato (*Lycopersicon esculentum* Mill). *International Journal of Current Microbiology and Applied Sciences*, 7(2), 2542-2545. <https://doi.org/10.20546/ijcmas.2018.702.309>
5. Muliarta, I. N. (2019, October). A study on rice field farmer implementation of rice straw composting. In *IOP Conference Series: Earth and Environmental Science*, 343(1), p. 012001. IOP Publishing. <https://doi.org/10.1088/1755-1315/343/1/012001>
6. Zhao, H., Yu, H., Yuan, X., Piao, R., Li, H., Wang, X., & Cui, Z. (2014). Degradation of lignocelluloses in rice straw by BMC-9, a composite microbial system. *Journal of Microbiology and Biotechnology*, 24(5), 585-591. <https://doi.org/10.4014/jmb.1310.10089>
7. Jain, M. S., Daga, M., & Kalamdhad, A. S., (2019). Variation in the key indicators during composting of municipal solid organic wastes. *Sustainable Environment Research*, 29(1), 1-8. <https://doi.org/10.1186/s42834-019-0012-9>
8. Chandna, P., Nain, L., Singh, S., & Kuhad, R. C. (2013). Assessment of bacterial diversity during composting of agricultural byproducts. *BMC Microbiology*, 13(1), 1-14. <https://doi.org/10.1186/1471-2180-13-99>
9. Rastogi, M., Nandal, M., & Khosla, B. (2020). Microbes as vital additives for solid waste composting. *Heliyon*, 6(2), e03343. <https://doi.org/10.1016/j.heliyon.2020.e03343>

10. Suliasih, & Widawati, S. (2018, November). The effect of biofertilizer combined with organic or inorganic fertilizer on growth of *Caesalpinia pulcherrima* and bacterial population in soil. In *IOP Conference Series: Earth and Environmental Science*, 166(1), p. 012024. IOP Publishing. <https://doi.org/10.1088/1755-1315/166/1/012024>
11. Kalbani, F. O. S. A., Salem, M. A., Cheruth, A. J., Kurup, S. S., & Senthilkumar, A. (2016). Effect of some organic fertilizers on growth, yield and quality of tomato (*Solanum lycopersicum*). *International Letters of Natural Sciences*, 53, 1-9. <https://doi.org/10.18052/www.scipress.com/ILN S.53.1>
12. Mulyani, O., Trinurani, E., Sudirja, R., & JoyOviyanti, B. (2017). The effect of bio-fertilizer on soil chemical properties of sugarcane in Purwadadi Subang. *KnE Life Sciences*, 2(6), 164-171. <https://doi.org/10.18502/klis.v2i6.1035>
13. Harman, G. E. (2006). Overview of mechanisms and uses of *Trichoderma* spp. *Phytopathology*, 96(2), 190-194. <https://doi.org/10.1094/PHYTO-96-0190>
14. Sawadogo, J., Ouédraogo, W. B. M., Coulibaly, P. J. A., Savadogo, C. A., Kaboré, A., & Legma, J. B. (2019). Étude comparative de la qualité de trois amendements organo-biologiques sur la production de tomate à Soala dans le CentreOuest du Burkina Faso [Comparative study of the quality of three organo-biological amendments on tomato production in Soala in the Center-West of Burkina Faso], *Science et Technique*, 38(1), 35-50.
15. Pindi, P. K., & Satyanarayana, S. D. V. (2012). Liquid microbial consortium- A potential tool for sustainable soil health. *Journal of Biofertilizer and Biopesticides*, 3(4), 2-4. <https://doi.org/10.4172/2155-6202.1000124>
16. Anwer, M. A., & Khan, M. R. (2013). *Aspergillus niger* as tomato fruit (*Lycopersicum esculentum* Mill.) quality enhancer and plant health promoter. *Journal of Postharvest Technology*, 1(1), 036-051.

JOURNAL OF SMART SCIENCE AND TECHNOLOGY



Editorial Board

Chief Editor

Khong Heng Yen, Universiti Teknologi MARA, Malaysia

Co-Chief Editor

Rudy Tawie, Universiti Teknologi MARA, Malaysia

Managing Editor

Robin Chang Yee Hui, Universiti Teknologi MARA, Malaysia

Co-Managing Editor

Leong Siow Hoo, Universiti Teknologi MARA, Malaysia

Advisory Board

Jamil Hamali, Universiti Teknologi MARA, Malaysia

Haeng-Ki Lee, KAIST, South Korea

Yoshinori Asakawa, Tokushima Bunri University, Japan

Editors

Mohamad Rusop Mahmood, Universiti Teknologi MARA, Malaysia

Yap Bee Wah, Universiti Teknologi MARA, Malaysia

Yana M. Syah, Institut Teknologi Bandung, Indonesia

Zhang Lei, Anhui Medical University, China

Bo Xia, Queensland University of Technology, Australia

Jesús del Barrio Lasheras, Universidad de Zaragoza, Spain

Wongi S. NA, Seoul National University of Science & Technology, South Korea

Hammad R. Khalid, King Fahd University of Petroleum & Minerals, Saudi Arabia

Siva K. Balasundram, Universiti Putra Malaysia, Malaysia

Sim Siong Fong, Universiti Sarawak Malaysia, Malaysia

Mohamad Hafiz Mamat, Universiti Teknologi MARA, Malaysia

Yupiter Harangan Prasada Manurung, Universiti Teknologi MARA, Malaysia

Ahmad Faiz Abdul Rashid, Universiti Teknologi MARA, Malaysia

Cindy Tan Soo Yun, Universiti Teknologi MARA, Malaysia

Harikesh Bahadur Singh, GLA University, India

Associate Editors

Asmah Alia Mohamad Bohari, Universiti Teknologi MARA, Malaysia

Clotilda Petrus, Universiti Teknologi MARA, Malaysia

Liew Fui Kiew, Universiti Teknologi MARA, Malaysia

Wan Fairos Wan Yaacob, Universiti Teknologi MARA, Malaysia

Academic Editors

Rudy Tawie

Rafidah Husen

Robin Chang Yee Hui

Wan Fairos Wan Yaacob

Chew Fei Ha

Ahmad Faiz Abdul Rashid

Copy Editor

Liew Chin Ying

Secretariat

Nyotia Nyokat

Yanti Yana Binti Halid

Widyani Darham

Rumaizah Haji Che Md Nor

Imelia Laura Daneil

Kimberley Lau Yih Long

Jacqueline Susan Rijeng @ Rejeng

Syah Runniza Ahmad Bakri

Hemyza Budin

Siti Faridah Kamaruddin

Webmaster

Lee Beng Yong

Cover Designer

Clement Jimel

WhatsApp

SISTER | Layanan_bkd

Journal of Smart Science and Technology

Editorial Policies | Journal of Smart Science and Technology

jsst.uitm.edu.my/index.php/jsst/ethics

Journal of Smart Science and Technology

Current Archives About

Search

Home / Editorial Policies

Editorial Policies

Ethical Guide for Editor

Editor should consider each manuscript unbiasedly.

Editor may reject a manuscript without review if it is incomplete or outside the journal's scope.

Editor must keep the confidentiality of all the information of a submission.

Editors should not involve in the review process of any self-authored manuscript submitted to the journal.

Editor should not use any material in a submission for other purposes without the consent of the authors.

Editor should make all reasonable efforts to process submissions on time.

Ethical Guide for Reviewers

Reviewer should understand the review policy of the journal.

Reviewer should declare any potential conflict of interest.

Reviewer should assess the manuscript objectively based on scientific quality. Avoid personal criticism of the author, and defamatory remarks.

Reviewer should express their views clearly with supporting arguments and references.

Reviewer should not use any material in a submission for other purposes.

Reviewer must keep the confidentiality of all the information of a submission.

Reviewer must keep the confidentiality of communication with the editor(s).

Reviewer must contact the editor if he/she wants to pass a review invitation onto another reviewer.

Peer Review Process

This journal operates a double-blind review process. The process is as follows:

1. The journal/ The publisher receives a manuscript.

2. The manuscript will be screened for plagiarism using Original. Only manuscripts with a similarity index of not more than 20% will be processed further.

3. An editor will be assigned to assess the manuscript for completeness and scope to decide whether it is suitable for review. In the case if an editor is on the author's list or has any conflict of interest, this editor will be excluded from handling and making decision on the manuscripts.

4. The handling editor identifies a minimum of two reviewers to assess the scientific quality of the manuscript. The review process should be completed within four (4) weeks.

5. The reviewers provide written reviews and suggest a recommendation for publication, which are either accept, reject, or revise.

6. The handling editor considers but not bound by the review reports for final decision regarding acceptance, rejection, or revision of the manuscript.

7. The corresponding author receives the reviewed report and the editorial decision.

8. If revision is required, the authors will be given 2 weeks (minor revision) or 8 weeks (major revision) to revise the manuscript.

9. The authors decide whether to revise and resubmit or withdraw.

10. If the manuscript is accepted, it moves into production for publication.

Platform & workflow by OJS / PKP

28°C

Hujan akan berh...

Search

05:04

23/02/2023