MANUSCRIPT HISTORY Published in Research on Crops (Q3)

Publisher: Gaurav Society of Agricultural Research Information Centre

https://www.scopus.com/sourceid/19900191751

Eile Edit View History Bookmarks Iools Help				3 × ~			
← → C ○ A https://www.scopus.com/sourceid/19900191751 ☆ Most Visited ♀ Getting Started			\$	⊚ ≡			
Scopus Preview Q Author Se Source details	earch Sources	⑦ ፹ Creat	e account Sign i	1			
Research on Crops Scopus coverage years: from 2008 to Present	0						
Publisher: Gaurav Society of Agricultural Research Information Centre SIR 2021 ISSN: 0972-3226 Sigr 2021 Subject area: (Agricultural and Biological Sciences: Agrinomy and Crop Science) (Agricultural and Biological Sciences: Soil Science) Sigr 2021 Source type: Journal Store type: Journal Sigr 2021 View all documents > Set document alert Save to source list Sigr 2021							
i Improved CiteScore methodology CiteScore 2021 counts the citations received in 2018-2021 to acticles, reviews, conference papers, book chapters and data			×	Ŷ			
🖽 🔎 Type here to search 🛛 🗄 🗟 💼 🚾 💽 👼 🚺 🚺		🕂 Hujan sekarang 🗸	へ (j) (12/21/20 12/21/20	22			

https://www.scimagojr.com/journalsearch.php?q=19900191751&tip=sid&clean=0

Eile Edit View History	Bookmarks Tools Help								_	5	/
Scopus preview	w - Scopus - Rese × 🔤 Research on Crops	×	Research on Crops	× +							,
$\leftrightarrow \rightarrow G$	O A https://www.scimagojr.	.com/jouri	nalsearch.php?q=19900191	751&tip=sid&clean=0				E \$. ⊡	=
🌣 Most Visited 💊 Getti	ing Started										^
						also	developed by scimago:		JTIONS RA	NKINGS	
SJR	Scimago Journal & Country Rank	<					Enter Journal	Title, ISSN or Publisher Name		Q	Ľ
		(and a	Jaumal Daphings	Country Depkings	Via Teele	Links	About Up				

Research on Crops

-

COUNTRY India Universities and research institutions in India	SUBJECT AREA AND CATEGORY Agricultural and Biological Sciences Agronomy and Crop Science Soil Science	PUBLISHER Gaurav Society of Agricultural Research Information Centre	H-INDEX 15
	ISSN	COVERAGE	
Type here to search	i 🖾 💽 🖬 🔟 🚺	1	+ Chillepage

https://indianjournals.com/ijor.aspx?target=ijor:rcr&type=for authors

File Edit View History Bookmarks Iools Help	-	Ø	\times						
😻 🚾 Scopus preview - Scopus - Rese × 📴 Research on Crops × 📴 Research on Crops × 🛄 Indianjournals.com//jor.aspx?ta × +			\sim						
← → C O A https://indianjournals.com//jor.aspx?target=ijor.rcr&type=for_authors		\bigtriangledown	\equiv						
The Most Visited 💊 Getting Started									
	(180.242.215.108)		^						
Andrian John Tars. Confin	•								
Home About us My Profile Registration Products Article Submission Usage Statistics Price List 2023 Contact Us Tutorial Login	n/Register								
Email id Log In	J								
ROC Author Guidelines									
Journal Home Carrent Issue Manuscripts: Redistration Subscripts									
Editorial Board Ams & Scope Author Guidelines	The full-length research papers and review articles written in English are welcome for publication in the 'Research on Crops' journal (Print ISSN 0972-3226; Online ISSN 2348-7542). The abbreviated title of the journal is 'Res. Crop.'.								
Ethics & Aims and Scope: Maipractice News & Events									
Subscribe TOC Nerts The "Research on Crops' is a peer-reviewed international journal publishing original research on basic and applied aspects of agronomy, soils and crop sciences including agricultural water ma agro-climatology, agroforestry, crop production, crop protection, cropping systems, genetics & plant breeding, horticulture, plant biotechnology, plant nutrition, post-harvest management of crop science, soil management & tillage, vegetables, weed science, agricultural engineering, environmental science, nano-technology, food science and technology, agri-business, agricultural econ	anagement, ips, seed nomics, etc.								
FREE The full-length research papers should be typed in one column on A4 size page leaving 2.5 cm margins on all sides. The manuscripts should be typed in MS Word format using Times N Sample Issue pl. font with 1.5 space between lines. Normally, the research papers containing more than 10 typed pages excluding tables and figures are not considered for publication in this journal.	New Roman 12								
The tables/figures should not cover more than 30% space of the text. Trial Access The full-length research papers submitted for publication should not contain data older than 3 years on the date of submission.	The tables/figures should not cover more than 30% space of the text. Trial Access The full-length research papers submitted for publication should not contain data older than 3 years on the date of submission.								
The papers written on the basis of one season/year field research are normally not considered for publication in 'Research on Crops' journal.	The papers written on the basis of one season/year field research are normally not considered for publication in 'Research on Crops' journal.								
The authors who are interested to publish field research in 'Research on Crops' journal should write their papers on the basis of minimum two seasons/years field research data.									
The papers written on the basis of one season/year research conducted under laboratory/controlled conditions can be considered for publication in 'Research on Crops' journal.									
🕂 🖉 Type here to search 🖄 🛱 🗟 🛱 😾 💽 🧮 💆 🚺 🏟	로 다) ENG 2:21 12/21	PM [/2022	\Box						



Manuscript submission : 3 Oktober 2022

COVER LETTER

To Research on Crops

Dear Editor,

I would like to send an original article entitled: "**The Role of Biochar Amendments in Improving the Properties of Acid Sulphate Soil**" for Research on Crops to consider. I confirm that this work is genuine and has not been published elsewhere, nor is it considered for publication elsewhere. We believe and hope that this manuscript is worthy of publication by Research on Crops. We are interested in publishing articles in this journal because it has an excellent reputation, so it is a matter of pride if published in Research on Crops. Here I attach the manuscript.

Thank you Best regards,

Agusalim Masulili Panca Bhakti University, Pontianak, West Kalimantan, Indonesia

The Role of Biochar Amendments in Improving the Properties of Acid Sulphate Soil

AGUSALIM MASULILI^{1,*}, SUTIKARINI², RINI SURYANI³, IDA AYU SUCI⁴, ISMAIL ASTAR⁵, HARDI DOMINIKUS BANCIN⁶, AND PAIMAN⁷

¹Department of Agrotechnology, Faculty of Agriculture, Universitas Panca Bhakti, Pontianak 78113, West Kalimantan, Indonesia

²Department of Agrotechnology, Faculty of Agriculture, Universitas PGRI Yogyakarta, Yogyakarta

55182, Indonesia Author email: ¹agusalim@upb.ac.id, ²sutikarini@upb.ac.id, ³rini.suryani@upb.ac.id, ⁴idaayusuci@upb.ac.id, ⁵ismailastar@upb.ac.id, ⁶hdbancin14@upb.ac.id,

⁷⁾paiman@upy.ac.id

*Corresponding author: Agusalim Masulili, Tilp: +628125631353, fax : 056177442, email: agusalim@upb.ac.id

ABSTRACT

Acid sulphate soil has the potential for agricultural development. Nevertheless, this soil has various soil properties problems that can inhibit plant growth. The granting of amendments can improve the properties of the soil. The study aimed to know the role of rice husk biochar combined with amendments to *Chromolaena odorata*, rice straw, and rice husk ash to improve the properties of acid sulphate soil. The study was conducted in a greenhouse and arranged in a complete randomized design (CRD) with three replications. Biochar amendments consisted of nine kinds, namely: without amendments (control), *Chromolaena odorata* (10 t/ha), rice straw (10 t/ha), rice husk biochar (1

ash (10 t/ha), *Chromolaena odorata* (10 t/ha) + rice husk biochar (5 t/ha), rice straw (10 t/ha) + rice husk biochar (5 t/ha), *Chromolaena odorata* (10 t/ha) + rice husk ash (5 t/ha), and rice straw (10 t/ha) + rice husk ash (5 t/ha). The results showed that the amendment of rice husk biochar combined with amendments to *Chromolaena odorata* and rice straw, respectively, had a good effect on improving soil physical properties, namely total soil pores, reducing soil content weight and soil strength. Its effect on soil chemical properties is that it can increase pH, C-organic, P-available, cation exchange capacity (CEC), and lower Al-dd and Fe soluble. The research findings show that applying biochar and organic amendments to *Chromolaena odorata* and rice straw can potentially improve the properties of acid sulphate soil. In future research, we recommend increasing the carrying capacity of acid sulphate soil against plant growth, and it can be done by improving soil properties through the organic amendments application of *Chromolaena odorata* or straw enriched with rice husk biochar.

Key words: *Chromolaena odorata*, rice husk ash, rice husk biochar, rice straw, soil properties **Running headline:** Improving the properties of acid sulphate soil

INTRODUCTION

Acidic sulphate soils are the common name for soils and sediments containing iron sulfide, namely pyrite (Das and Das, 2015). In agricultural development, efforts to reclaim acid sulphate soils do not always result in soil productivity since improvements in soil properties after reclamation are not achieved. So far, efforts have been encouraged to overcome existing obstacles after reclamation, including inorganic fertilizers (urea, TSP, KCl) and liming (dolomite). However, rice crop productivity in this field has not been optimal.

Inorganic fertilizers increase over time and can even decrease land quality if inputs are high and intensive. Liming can only cope with short-term pH and is often done repeatedly. As a result, high costs are needed, which becomes a burden for farmers and will threaten the sustainability of land productivity. Therefore, it is necessary to manage land based on land amendments that can improve the nature of sustainable soils.

Reclamation can result in changes in soil properties and is related to the potential for pyrite oxidation, affecting the increase in soil packaging and the solubility of ions such as Al, Fe, Mn, and sulphate. On the other hand, this soil has a high clay content which affects the increased weight of the contents. Therefore, the development of acid sulphate soil for agriculture after reclamation requires proper management of soil amendments to achieve the formation of steady cultivated land and sustainable farming. Das and Das (2015) suggested that acid sulphate soil having a high pH can inhibit plant growth. In addition, when this soil is drained, it can affect an increase in sulfuric acid and the release of iron, aluminum, and other heavy metals. Shamshuddin *et al.* (2014), stated that plants grown in acidic sulphate soils could not grow well due to low pH and high Al content.

Biochar is one of the soil amendment materials with the long-term potential to maintain soil fertility. The results of experts' investigations into the phenomenon of terra preta show that soil fertility is stable in the Amazon basin due to the presence of black carbon, the result of the management of the Amerindian nation 500–2,500 years ago. This carbon black is nothing but biochar immersed in the soil. Murtaza *et al.* (2021) found that 4.0 t/ha cotton biochar improves physical, chemical, soil biological, and soil fertility properties and peanut productivity in Alfisol. Zhang *et al.* (2022) have proven that biochar provides effects such as liming and providing nutrients that lead to soil improvement. Kocsis *et al.* (2022) stated that biochar has a high porosity and can improve the physicochemical and biological properties of the soil. In line, Sovova *et al.* (2021) found biochar had a good influence on improving the soil properties of regosol. Further afield, Das *et al.* (2021) suggest the application of biochar to the soil can provide agronomic benefits, support crop yield improvement, and improve soil quality and health. In addition, biochar has been shown to act as a fertilizer and soil conditioner.

Nutrient retention and higher nutrient availability were found in the soil after biochar addition, associated with higher CEC, surface acreage, and direct nutrient addition (Glaser *et al.*, 2002). In line with opinions, Murtaza *et al.* (2021) suggested that biochar is a recalcitrant carbon product of

biomass produced through the pyrolysis process. The recalcitrant properties of biochar are distinctive and important to the long-term improvement of soil properties through the effects of their residues that can persist for a long time in the soil. Beusch (2021), biochar can contribute to soil fertility and crop productivity, and Kalu *et al.* (2021) increase water availability and lower soil BD.

Applying biochar can improve soil properties and long-term crop productivity compared to other organic amendments. In other organic amendments, research results by Suri and Yudono (2020) showed that giving *Cromolaena odorata* as compost can increase plants' nutrition and nutrient uptake. Agustina *et al.* (2019) argued that using *Chromolaena odorata* and rice straw can maintain soil quality. Mahmoud *et al.* (2009) also showed rice straw compost could improve soil quality. The same was discovered by Gewaily (2019), that straw compost can increase potassium availability.

Furthermore, the results of the study by Yin *et al.* (2022) showed the weight of the soil contents decreased, and the water-holding capacity and total porosity increased in the soil with rising rice husk ash. Adding rice husk ash neutralizes soil acidity and increases plant nutrient availability. Research by Lestari and Rachmawati (2020) showed that applying rice husk ash (4 t/ha) can increase rice growth and reduce salinity stress.

Previously, several studies on the effect of biochar singularly have been carried out, including on the P-availability (Zhang *et al.*, 2022), improvement of soil physical properties (Aslam *et al.*, 2014), chemical properties of the soil (Bista *et al.*, 2019), and some soil characteristics (Yadav *et al.*, 2018), soil aggregate stability (Ebido *et al.*, 2021), and to the development of rice roots (Kartika *et al.*, 2021). However, no research has been found on how the role of biochar was given together with other organic amendments in acid sulphate soil. Therefore, it is necessary to study how the effect of biochar is given together with organic amendments to *Chromolaena odorata* and rice straw. This research is expected to contribute to improving the nature of acid sulphate soil in West Kalimantan, which has been one of the types of soil widely used as an agricultural activity for rice crops. In connection with this hah, this study aimed to know the role of rice husk biochar combined with amendments to *Chromolaena odorata*, rice straw, and rice husk ash in improving the properties of acid sulphate soil.

MATERIALS AND METHODS

Study Area

This research was conducted from September to November 2021. The research site was carried out at the Laboratory of the Faculty of Agriculture, Universitas Panca Bhakti, Pontianak, West Kalimantan Province, Indonesia. The height of the study site is one m from above sea level, with an average temperature and humidity of 27.6 °C and 82.8%, respectively. The research position is located at a latitude of 2°05' N–3° 05' S and longitude of 108°30' E–144°10' W.

Experimental Design

The study was conducted in a greenhouse and arranged in a complete randomized design (CRD) with three replications. Biochar amendments consisted of nine kinds, namely: without amendments (control), *Chromolaena odorata* (10 t/ha), rice straw (10 t/ha), rice husk biochar (10 t/ha), rice husk ash (10 t/ha), *Chromolaena odorata* (10 t/ha) + rice husk biochar (5 t/ha), rice straw (10 t/ha) + rice husk biochar (5 t/ha), rice straw (10 t/ha) + rice husk biochar (5 t/ha). In all, 27 polybags are needed.

Research Procedures

The material in this study used acid sulphate soil taken on the experimental land of the West Kalimantan, Sungai Kakap, Agricultural Technology Assessment Center, and land farmers' work on rice crops. This area included the development of swampland agriculture which has built irrigation canals since 1980 and was currently designated as an Integrated Farmer Business Area. Soil picking

was carried out randomly at a soil depth of 0-20 cm, then mixed, air-dried, and cleaned. Then put in polybags was as much as 8 kg.

Biochar was made by taking rice husk from grain harvested from rice plants grown directly in the acid sulphate soil of the Kakap River, West Kalimantan, Indonesia. First, the rice mill was taken the grain to obtain rice husk. Furthermore, the rice husk was placed on the pyrolysis device to be processed into biochar by incomplete combustion (limited air) in a reactor made of Pertamina drums.

Organic amendments to *Chromolaena odorata* were taken from around the field, then chopped to a size of 1-2 cm. Rice straw was taken from waste harvested rice grown and grown by farmers in acid sulphate soil and chopped within 1-2 cm. Rice husks were taken from waste from rice milling, then used as ash through burning in open areas to produce grayish-white ash.

Prepared acid sulphate soil introduced into 27 polybags until it reached 8 kg. Further, the dose of soil amendment was calculated based on the soil weight in the polybag area, mixed with the soil in each polybag to a soil depth of 20 cm, after which it was incubated with moisture content until it was closed to the field capacity for 30 days.

At 30 days after incubation, from each study unit, soil sampling of about 50 g was undisturbed to determine the weight of the soil moisture content accident. In addition, a disturbed soil sample of 100 g was taken from each research unit for analysis of chemical properties.

Observation parameters

The soil bulk density (BD) was determined by the clod method described by Blake & Harke. The total pore was calculated from the soil moisture content (v/v) at the matrix potential of 0 kPa. The available groundwater was estimated by reducing the soil moisture content in the matrix potential by -33 kPa (field capacity) with the soil moisture content in the matrix potential, which was -15 Mpa (wilting point). The soil moisture content at this potential matrix was determined using a pressure plate device. The strength of the soil was measured with a hand penetrometer (Daikie) at a soil depth of 15 cm.

The pH level was measured in a soil solution at a ratio of 1:2.5 (with deionized water), using a pH meter (Jenway 3305). Walkley and Black's wet oxidation method was used to determining organic C levels (Soil Survey Laboratory Staff). Total N levels were measured by the Kjeldhal (Bremner and Mulvaney) method. $Al_3 + and Fe_2 + were$ extracted with 1 M KCl (Barnhisel and Bertsch). CEC was extracted with 1 M NH₄Oac (buffer at pH 7.0), and the concentration of alkaline cations was measured using AAS (Shimadzu), P-available with Bray I.

Statistical Analysis

Statistical analysis was carried out on each observation parameter data to determine the effect of the amendments on the nature of acid sulphate soil. The data obtained were analyzed by analysis of variance (ANOVA) at 5% significant level. In addition, the test of least significant difference (LSD) at 5% significant level was used to determine the average value between treatments.

RESULTS AND DISCUSSION

Effect of various soil amendments on some physical soil properties

Applying biochar improved the soil's physical properties (Aslam *et al.*, 2014). Applying biochar and organic amendments influenced the changes in some of the observed soil physical properties. The statistical test is shown in Table 1. The soil BD was lower on all amendments granted. For example, in soils fed a single rice husk biochar (10 t/ha), soil BD decreased from 1.27 mg/m³ in the control soil to 1.15 mg/cm³. However, not significantly different from soils was given straw (10 t/ha) and a combination of straw + biochar, *Chromolaena odorata* + rice husk ash, and straw + rice husk

ash. Furthermore, in the combination of *Cromolaena odorata* (10 t/ha) + rice husk biochar (5 t/ha), a BD of 1.13 mg/m^3 was obtained and was the lowest soil BD. Still, it was not significantly different from the soil, given a combination of straw (10 t/ha) + rice husk biochar 5 (t/ha) produced soil BD of 1.14 mg/m^3 . These results indicate an improvement in the physical soil properties if biochar was given together with organic amendments to *Chromolaena odorata*.

	BD	Total	Penetration resistance (N/cm ²)				
Soll amendments	(mg/m^3)	pores (%)	pF 0	pF 2	pF 2.5		
Control	1.24 e	44.43 a	36.67 b	310.00 e	500.00 d		
C. odorata (10 t/ha)	1.16 c	52.17 e	20.00 a	230.00 abc	403.33 bc		
Rice straw (10 t/ha)	1.17 bc	53.27 f	16.67 a	243.33 bc	393.33 b		
Rice husk biochar (10 t/ha)	1.15 b	54.21 g	14.33 a	223.33 ab	390.00 b		
Rice husk ash (10 t/ha)	1.19 d	47.30 b	20.00 a	270.00 d	403.33 bc		
C. odorata (10 t/ha) +	1.13 a	56.73 i	10.00 a	220.00 a	340.00 a		
biochar (5 t/ha)							
Rice straw (10 t/ha) + rice	1.14 ab	55.57 h	13.33 a	223.00 ab	360.00 a		
husk biochar (5 t/ha)							
C. odorata (10 t/ha) + rice	1.15 b	48.90 c	16.67 a	240.00 bc	420.00 c		
husk ash (5 t/ha)							
Rice straw (10 t/ha) + rice	1.16 bc	50.30 d	20.00 a	246.67 с	403.33 bc		
husk ash (5 t/ha)							

Table 1. Effect of rice husk biochar with soil amendments on some physical properties of acid sulphate soil at one month after incubation

Remarks: The average numbers followed by the same letter in the same column show no significant difference in the LSD test at 5% significant level.

Applying various biochar amendments and their combinations increased the soil's total pore. The statistical test results are shown in Table 1. There was a significant difference in the total pore of the soil between the various amendments. Control soils had the lowest total pore (44.43%), contrasting with soils that provided various amendments. On soils fed with rice husk biochar (10 t/ha), the total soil pore increased to 54.21%, significantly different from other single amendments. Furthermore, a combination of *Cromolaena odorata* (10 t/ha) + rice husk biochar (5 t/ha) caused the total pore to increase to 56.73% and was the highest total pore.

The application of various amendments, both single and combined, affected the decrease in soil strength as measured as soil penetration resistance (N/cm²) at conditions of pF 0, pF 2, and pF 2.5, where the higher the soil matric suction (pF), the higher the soil penetration resistance. The results of statistical tests (Table 1) show that under saturated conditions (pF 0), the various soil amendments do not offer a noticeable difference. In pF 2 shows that the application of rice husk biochar (10 t/ha) can reduce soil penetration resistance from 310.00 N/cm² (control soil) to 223.33 N/cm². No significant difference between soils fed with *Chromolaena odorata* and rice straw or a combination of straw (10 t/ha) + rice husk biochar (5 t/ha). Furthermore, soils given a combination of *Cromolaena odorata* (10 t/ha) + biochar (5 t/ha) obtained the lowest penetration resistance (220.00 N/cm²). Still, it was not significantly different from soils have given *Chromolaena odorata* (10 t/ha), rice husk biochar (10 t/ha) + biochar (5 t/ha) + biochar (5 t/ha) + biochar (5 t/ha) biochar (5 t/ha) + biochar (5 t/ha).

Similarly, at pF 2.5, the application of rice husk biochar (10 t/ha) was single. It could reduce the value of soil penetration resistance from 500.00 N/cm² on control soil to 390.00 N/cm². Still, it was not significantly different from soils have given *Chromolaena odorata* (10 t/ha), straw (10 t/ha), rice husk ash (10 t/ha), and a combination of rice straw (10 t/ha) + rice husk ash (5 t/ha). Furthermore, the combination of *Cromolaena odorata* (10 t/ha) + rice husk biochar (5 t/ha) achieved the lowest penetration resistance (340.00 N/cm²). Still, it was not significantly different from the combination

of rice straw (10 t/ha) + biochar (5 t/ha) (360.00 N/cm²). It indicated that rice husk biochar had the potential to control soil strength. In line with Das *et al.* (2021), applying biochar into the soil could reduce soil strength.

Effect of rice husk biochar on some soil chemical properties

The application of several soil amendments and their combinations could influence some chemical properties of acid sulphate soil. The results of statistical tests in Table 2 show that soil reaction (pH) of the soil increased in all treatments given compared to controls. For example, in the application of rice husk biochar showed an increase in pH from 3.36 (control soil) to 4.40 on soils given biochar (10 t/ha) but did not significantly different from other single amendments. Furthermore, the combined application of *Cromolaena odorata* (10 t/ha) + biochar (5 t/ha) obtained the highest increase in soil pH (4.58), in marked contrast to soils given rice husk ash but not significantly different from single amendments or other combinations.

Applying single and combined soil amendments affected to increase in soil C-organics. The statistical test results in Table 2 show that the control soil had the lowest C-organic (1.94%), in stark contrast to other amended soils, except that the soils have given rice husk ash (10 t/ha) was not significantly different. The soil given a single amendment of rice straw (10 t/ha) obtained 4.58% C-organic, in stark contrast to the control soil and the soil given rice husk ash, but not significantly different from other soil amendments. Similarly, soils given a combination of *Chromolaena odorata* (10 t/ha) + biochar husk (5 t/ha) achieved the highest soil C-organic (4.93%) in marked contrast to control soils and soils fed with rice husk ash but did not significantly different from other soil amendments.

Soil amendments	pH H ₂ O	C- organic (%)	P- available (%)	Al-dd (me/100 g)	Soluble Fe (%)	CEC (me/100 g)
Control	3.36 a	1.94 a	0.21 a	3.84 e	3.61 d	6.64 a
C. odorata (10 t/ha)	4.06 bc	4.22 b	0.29 bc	3.31 c	3.28 c	7.15 ab
Rice straw (10 t/ha)	4.28 bc	4.58 b	0.30 bc	3.42 cd	3.34 cd	7.32 abc
Rice husk biochar (10 t/ha)	4.40 bc	4.09 b	0.32 cd	3.16 b	3.10 abc	8.03 c
Rice husk ash (10 t/ha)	3.98 b	2.78 a	0.28 b	3.51 d	3.34 cd	7.76 bc
C. odorata (10 t/ha) + biochar (5 t/ha)	4.58 c	4.93 b	0.34 d	3.02 a	2.94 ab	10.04 e
Rice straw (10 t/ha) + rice husk biochar (5 t/ha)	4.48 bc	4.75 b	0.33 d	3.08 ab	2.91 a	9.67 de
<i>C. odorata</i> (10 t/ha) + rice husk ash (5 t/ha)	4.26 bc	4.70 b	0.31 cd	3.37 c	3.25 c	9.17 d
Rice straw (10 t/ha) + rice husk ash (5 t/ha)	4.26 bc	4.73 b	0.31 cd	3.47 d	3.22 bc	9.20 d

Table 2. Effect of rice husk biochar with soil amendments on some chemical properties of acid sulphate soil at one month after incubation

Remarks: The average numbers followed by the same letter in the same column show no significant difference in the LSD test at 5% significant level.

Applying various amendments and combinations to acid sulphate soil generally affected the increase in P-available. The results of statistical tests (Table 2) showed that control soils had the lowest P-available content (0.21%), significantly different from soils given various amendments. In soils fed with a rice husk biochar (10 t/ha), the P-available increased to 0.32%. Still, it was not significantly different from other combined or single-amended soils, except with soils that were given rice husk ash with a lower P-available (0.27%). The highest P-available was obtained on soils given a combination of *Chromolaena odorata* (10 t/ha) + biochar (5 t/ha) (0.34%), in stark contrast to control soils and those topped with rice husk ash but not significantly different from single amendments or other combinations.

The decrease in Al-dd and Fe-dissolve in acid sulphate soil also occurs due to applying various soil amendments and their combinations. Statistical tests (Table 2) show that the highest Al-dd was obtained on control soils (3.84%), further decreasing and differing markedly from soils given various single and combined amendments. On single-amendment soils of rice husk biochar (10 t/ha), Al-dd decreased to 3.16%, significantly different from other single-amended soils. Furthermore, the soil given the combination of *Chromolaena odorata* (10 t/ha) + rice husk biochar (5 t/ha) obtained the lowest Al-dd content (3.02%). It was not significantly different from the soil given a combination of rice straw (10 t/ha) + biochar (5 t/ha), but it was significantly different from other amended soils. In soluble Fe content, the application of rice husk biochar (10 t/ha) could reduce soluble Fe from 3.61% on control soils to 3.10% and was significantly different from other single amendments. And when the soil was given a combination of rice straw (10 t/ha) + rice husk biochar (5 t/ha) caused the lowest decrease in Al-dd (2.91%). Still, it was not significantly different from soil that was given a combination of *Chromolaena odorata* (10 t/ha) + rice husk biochar (5 t/ha).

The CEC of land has also increased due to the application of various land amendments and their combinations. The statistical test is shown in Table 2 show that the control soil had the lowest CEC (6.64 Cmol/kg), not significantly different from soils fed with *Chromolaena odorata* and rice straw. On the other hand, on soils provided with rice husk biochar (10 t/ha), the soil CEC increased to 8.03 Cmol/kg, but it was not significantly different from soil given rice straw and rice husk ash.

In line with the results that have been put forward, Yamato *et al.* (2006) found that the application of biochar from *Acacia mangium* bark into the soil in Sumatra led to changes in soil chemical properties through increased pH, N-total and P-available, CEC, and lowered interchangeable Al. Liang *et al.* (2006) found that two mechanisms could create a larger CEC. First, a higher charged density per unit of surface acreage means a higher degree of oxidation of soil organic matter. Second, the presence of a higher surface area for cation absorption or the combined effect of both. Glaser *et al.* (2002) posit that oxidation of aromatic C and carboxyl formation are the main reasons for the high CEC. Liang *et al.* (2006) also found that the CEC per unit C was higher, and the load density was higher in carbon-black-rich Anthrosol compared to soils that were poor in carbon black. In addition, Anthrosol showed a higher surface acreage due to its higher carbon black concentration.

From the results, the great potential of rice husk biochar against improving soil properties increased when given together with the biomass *Chromolaena odorata*. It could happen because *Chromolaena odorata* contains higher humate and fulvic organic acids, so that it could encourage the complex mechanisms of organo-mineral organs and soil aggregation. In addition, *Chromolaena odorata* can highly suppress Al and Fe, thereby increasing the P availability in the soil due to the presence of organic acids released during decomposition. Research results by Suri and Yudono (2020) also found that using *Chromolaena odorata* compost could improve soil quality and nutrient uptake of lettuce plants.

Chromolaena odorata can contribute to the care of soil C-organic and physical properties due to a reasonably high polyphenol content that could inhibit the decomposition and mineralization of N organic matter, even though it had a low C/N ratio.

CONCLUSIONS

In conclusion, the amendment of rice husk biochar combined with amendments to *Chromolaena* odorata and rice straw improved soil physical properties, namely total soil pores, reducing soil

content weight and strength. Its impact on soil chemical properties was that it could increase pH, Corganic, P-available, CEC, lower Al-dd, and Fe soluble. Therefore, applying biochar and organic amendments to *Chromolaena odorata* and rice straw could improve the properties of acid sulphate soil. Consequently, we recommend increasing the carrying capacity of acid sulphate soil against plant growth. It can be done by enhancing soil properties through the organic amendments application of *Chromolaena odorata* or straw enriched with rice husk biochar.

ACKNOWLEDGEMENTS

We thank the Institute for Research and Community Service, Universitas Panca Bhakti, which has given support for research funds. In addition, we thank the Faculty of Agriculture, Universitas Panca Bhakti, which has provided loans for facilities in the form of equipment for research.

REFERENCES

- Agustina, R., Jumadi, R., Firmani, U. and Abdul Faisal, R. H. (2019). Effects of decomposition rate of *Chromolaena odorata* and straw rice in fresh and compost form to the growth and yield of rice. *IOP Conf Ser Earth Environ Sci.* 250(1): 1–5. <u>https://doi.org/10.1088/1755-1315/250/1/012078</u>
- Aslam, Z., M. Khalid, M. and Aon, M. (2014). Impact of biochar on soil physical properties. Sch J Agric Sci. 4(5): 280–84.
- Beusch, C. (2021). Biochar as a soil ameliorant: How biochar properties benefit soil fertility A review. J Geosci Environ Prot. 9(10): 28–46. https://doi.org/10.4236/gep.2021.910003
- Bista, P., Ghimire, R., Machado, S. and Pritchett, L. (2019). Biochar effects on soil properties and wheat biomass vary with fertility management. *Agronomy*. **9**(10): 1-10. https://doi.org/10.3390/agronomy9100623
- Das, S. K. and Das, S. K. (2015). Acid sulphate soil: Management strategy for soil health and productivity. *Pop. Kheti.* **3**(2), 3–8.
- Das, S., Mohanty, S., Sahu, G., Rana, M. and Pilli, K. (2021). Biochar: A sustainable approach for improving soil health and environment. In: Soil Erosion - Current Challenges and Future Perspectives in a Changing World, (April). https://doi.org/10.5772/intechopen.97136
- Ebido, N. E., Edeh, I. G., Unagwu, B. O., Nnadi, A. L., Ozongwu, O. V., Obalum, S. E. and Igwe, C. A. (2021). Rice-husk biochar effects on organic carbon, aggregate stability and nitrogen-fertility of coarse-textured Ultisols evaluated using *Celosia argentea* growth. *Sains Tanah*. 18(2): 177–87. https://doi.org/10.20961/stjssa.v18i2.56330
- Gewaily, E. (2019). Impact of compost rice straw and rice straw as organic fertilizer with potassium treatments on yield and some grain quality of Giza 179 rice variety. *J Plant Prod.* 10(2): 143– 151. https://doi.org/10.21608/jpp.2019.36244
- Glaser, B., Lehmann, J. and Zech, W. (2002). Ameliorating physical and chemical properties of highly weathered soils in the tropics with charcoal - A review. *Biol Fertil Soils*. 35(4): 219–30. <u>https://doi.org/10.1007/s00374-002-0466-4</u>
- Kalu, S., Simojoki, A., Karhu, K. and Tammeorg, P. (2021). Long-term effects of softwood biochar on soil physical properties, greenhouse gas emissions and crop nutrient uptake in two contrasting boreal soils. *Agric Ecosyst Environ.* 316: 107454. https://doi.org/10.1016/j.agee.2021.107454
- Kartika, K., Sakagami, J. I., Lakitan, B., Yabuta, S., Akagi, I., Widuri, L. I., Siaga, E., Iwanaga, H. and Nurrahma, A. H. I. (2021). Rice husk biochar effects on improving soil properties and root development in rice (*Oryza glaberrima* Steud.) exposed to drought stress during early reproductive stage. *AIMS Agric Food.* 6(2): 737–751. https://doi.org/10.3934/AGRFOOD.2021043
- Kocsis, T., Ringer, M. and Biró, B. (2022). Characteristics and applications of biochar in soil–plant systems: A short review of benefits and potential drawbacks. *Appl Sci.* **12**(8): 1–16. https://doi.org/10.3390/app12084051
- Lestari, M. F. and Rachmawati, D. (2020). The effect of rice husk ash on growth of rice (Oryza sativa

L. 'Sembada Merah') on salinity stress. *AIP Conf Proc.* **2260**: 1–6. https://doi.org/10.1063/5.0015860

- Liang, B., Lehmann, J., Solomon, D., Kinyangi, J., Grossman, J., O'Neill, B., Skjemstad, J. O., Thies, J., Luizão, F. J., Petersen, J. and Neves, E. G. (2006). Black carbon increases cation exchange capacity in soils. *Soil Sci Soc Am J*. **70**(5): 1719–30. https://doi.org/10.2136/sssaj2005.0383
- Mahmoud, E., Ibrahim, M., Robin, P., Akkal-Corfini, N. and El-Saka, M. (2009). Rice straw composting and its effect on soil properties. *Compost Sci Util.* **17**(3): 146–50. <u>https://doi.org/10.1080/1065657X.2009.10702415</u>
- Murtaza, G., Ahmed, Z., Usman, M., Tariq, W., Ullah, Z., Shareef, M., Iqbal, H., Waqas, M., Tariq, A., Wu, Y., Zhang, Z. and Ditta, A. (2021). Biochar induced modifications in soil properties and its impacts on crop growth and production. *J Plant Nutr.* 44(11): 1677–91. https://doi.org/10.1080/01904167.2021.1871746
- Shamshuddin, J., Elisa Azura, A., Shazana, M. A. R. S., Fauziah, C. I., Panhwar, Q. A. and Naher, U. A. (2014). Properties and management of acid sulfate soils in Southeast Asia for sustainable cultivation of rice, oil palm, and cocoa. Advances in Agronomy (1st ed., Vol. 124). Elsevier Inc. https://doi.org/10.1016/B978-0-12-800138-7.00003-6
- Sovova, S., Enev, V., Smilek, J., Kubikova, L., Trudicova, M., Hajzler, J. and Kalina, M. (2021). The effect of biochar application on soil properties and growth of the model plant Zea mays. *Ecocycles*. **7**(2): 46–54. https://doi.org/10.19040/ecocycles.v7i2.201
- Suri, A. M. and Yudono, P. (2020). Effects of *Chromolaena odorata* compost on soil and nutrient uptake of lettuce (*Lactuca sativa*). *Planta Trop J Agro Sci.* 8(1): 33–8. <u>https://doi.org/10.18196/pt.2020.111.33-38</u>
- Yadav, N. K., Kumar, V., Sharma, K. R., Choudhary, R. S., Butter, T. S., Singh, G., Kumar, M. and Kumar, R. (2018). Biochar and their impacts on soil properties and crop productivity: A review. *J Pharmacogn Phytochem Harpole*. 7(4): 49–54.
- Yamato, M., Okimori, Y., Wibowo, I. F., Anshori, S. and Ogawa, M. (2006). Effects of the application of charred bark of *Acacia mangium* on the yield of maize, cowpea and peanut, and soil chemical properties in South Sumatra, Indonesia. *Soil Sci Plant Nutr.* 52(4): 489–95. https://doi.org/10.1111/j.1747-0765.2006.00065.x
- Yin, M., Li, X., Liu, Q. and Tang, F. (2022). Rice husk ash addition to acid red soil improves the soil property and cotton seedling growth. *Sci Rep.* 12(1): 1–9. https://doi.org/10.1038/s41598-022-05199-7
- Zhang, Y., Chen, H., Xiang, J., Xiong, J., Wang, Y., Wang, Z. and Zhang, Y. (2022). Effect of ricestraw biochar application on the acquisition of rhizosphere phosphorus in acidified paddy Soil. *Agronomy*. 12: 1-13.

Online manuscript submission : 3 Oktober 2022

=	M Gmail		Q, is:important		× 莊	• Aktif • 💿	©	@UPF	
ail	/ Tulis		Dari 👻 Kapan saja 👻 A	da lampiran Kepada 👻 Belum dibaca Pen	elusuran lanjutan				
nat	 ☑ Kotak Masuk ☆ Berbintang ③ Ditunda 	5.340	🗌 👻 G :	Kotak Masuk Manuscript submission - Dear Dr. Vedpal	Singh Cief Editor Thank y	ou for your response. Best	regards, A	3 Okt	
aces Dr eet	Terkirim Toraf Ciutkan	4	 A noreply A Scholastica 	Kotak Masuk Konfirmasi Kode Pembayaran Tiket Pesaw Kotak Masuk [The Abstract] PRW highlights + Similarity	vat - [862801142] - Terima v Check preview - In this i	a kasih atas pesanan Anda! ssue: Peer Review Week hig	Tiket pesa ghlights an	1 Okt 30 Sep	
	 Penting Fo Terjadwal 		🔲 🚖 Dr. Muhammad Achiru.	Kotak Masuk [Teknotan] Editor Decision - Yth. sutikarin	ii SP, M.Sc: Terimakasih te	alah mengajukan naskah pa	da Teknot	26 Sep	
	😰 Semua Email 🛈 Spam		 	Kotak Masuk Everything you need to know about Peer	Review Week 2022 - Peer ak balek, menguii kesabar	Review Week 2022 is finall	y here, ru	20 Sep 14 Sep	
	III Sampah ▶ D Kategori		🗌 🏫 Dr. Muhammad Achiru.	(Kotak Masuk) [Teknotan] Keputusan Editor - Terimakasi	ih telah menyerahkan arti	kel anda yang berjudul, "PE	NGGUNA	14 Sep	
	 Kelola label + Buat label baru 			Kotak Masuk [MJFAS] Password Reset - Your new pass Kotak Masuk [MJFAS] Password Reset Confirmation - I	word: FHtfmT Reset my password: https	://mjfas.utm.my/index.php/r	mjfas/logi	11 Sep	
	Label	+	🗋 🛧 Kedaireka	Kotak Masuk [Matching Fund] Proposal Tidak Lolos Sel	eksi - Hai, Dr. Ir. Agusalim	Masulili, MP. Kami menguc	apkan teri	5 Sep	









Role of biochar amendments in improving the properties of acid sulphate soil AGUSALIM MASULILI^{1,*}, SUTIKARINI¹, RINI SURYANI¹, IDA AYU SUCI¹, ISMAIL ASTAR¹, HARDI DOMINIKUS BANCIN¹ AND PAIMAN²

¹Department of Agrotechnology, Faculty of Agriculture Panca Bhakti University, Pontianak 78113, West Kalimantan, Indonesia *(e-mail: <u>agusalim@upb.ac.id</u>) ²Department of Agrotechnology, Faculty of Agriculture, Universitas PGRI Yogyakarta, Yogyakarta 55182, Indonesia

ABSTRACT

Acid sulfate soils have potential for agricultural development. However, this soil has various properties problems that can inhibit plant growth. One way to improve soil properties is through the application of biochar amendments. The study aimed to know the role of rice husk biochar combined with amendments to *Chromolaena odorata*, rice straw, and rice husk ash to improve the properties of acid sulphate soil. The study was conducted in a greenhouse and arranged in a complete randomized design (CRD) with three replications. Biochar amendments consisted of nine kinds, namely: without amendments (control), *Chromolaena odorata* (10 t/ha), rice straw (10 t/ha), rice husk biochar (10 t/ha), rice husk biochar (5 t/ha), rice

straw (10 t/ha) + rice husk biochar (5 t/ha), *Chromolaena odorata* (10 t/ha) + rice husk ash (5 t/ha), and rice straw (10 t/ha) + rice husk ash (5 t/ha). The results showed that the amendment of rice husk biochar combined with amendments to *Chromolaena odorata* and rice straw, respectively, had a good effect on improving soil physical properties, namely total soil pores, reducing soil content weight and soil strength. Its effect on soil chemical properties is that it can increase pH, C-organic, P-available, cation exchange capacity (CEC), and lower Al-dd and Fe soluble. The research findings show that applying biochar and organic amendments to *Chromolaena odorata* and rice straw can potentially improve the properties of acid sulphate soil. In future research, we recommend increasing the carrying capacity of acid sulphate soil against plant growth, and it can be done by improving soil properties through the organic amendments application of *Chromolaena odorata* or straw enriched with rice husk biochar.

Key words: Chromolaena odorata, rice husk ash, rice husk biochar, rice straw, soil properties

INTRODUCTION

Acidic sulphate soils are the common name for soils and sediments containing iron sulfide, namely pyrite (Das and Das, 2015). In agricultural development, efforts to reclaim acid sulphate soils do not always result in soil productivity since improvements in soil properties after reclamation are not achieved. So far, efforts have been encouraged to overcome existing obstacles after reclamation, including inorganic fertilizers (urea, TSP, KCl) and liming (dolomite). However, rice crop productivity in this field has not been optimal.

Inorganic fertilizers increase over time and can even decrease land quality if inputs are high and intensive. Liming can only cope with short-term pH and is often done repeatedly. As a result, high costs are needed, which becomes a burden for farmers and will threaten the sustainability of land productivity. Therefore, it is necessary to manage land based on land amendments that can improve the nature of sustainable soils.

Reclamation can result in changes in soil properties and is related to the potential for pyrite oxidation, affecting the increase in soil packaging and the solubility of ions such as Al, Fe, Mn, and sulphate. On the other hand, this soil has a high clay content which affects the increased weight of the contents. Therefore, the development of acid sulphate soil for agriculture after reclamation requires proper management of soil amendments to achieve the formation of steady cultivated land and sustainable farming. Das and Das (2015) suggested that acid sulphate soil having a high pH can inhibit plant growth. In addition, when this soil is drained, it can affect an increase in sulfuric acid and the release of iron, aluminum, and other heavy metals. Shamshuddin *et al.* (2014), stated that plants grown in acidic sulphate soils could not grow well due to low pH and high Al content.

Biochar is one of the soil amendment materials with the long-term potential to maintain soil fertility. The results of experts' investigations into the phenomenon of terra preta show that soil

fertility is stable in the Amazon basin due to the presence of black carbon, the result of the management of the Amerindian nation 500–2,500 years ago. This carbon black is nothing but biochar immersed in the soil. Murtaza *et al.* (2021) found that 4.0 t/ha cotton biochar improves physical, chemical, soil biological, and soil fertility properties and peanut productivity in Alfisol. Zhang *et al.* (2022) have proven that biochar provides effects such as liming and providing nutrients that lead to soil improvement. Kocsis *et al.* (2022) stated that biochar has a high porosity and can improve the physicochemical and biological properties of the soil. In line, Sovova *et al.* (2021) found biochar had a good influence on improving the soil properties of regosol. Further afield, Das *et al.* (2021) suggest the application of biochar to the soil can provide agronomic benefits, support crop yield improvement, and improve soil quality and health. In addition, biochar has been shown to act as a fertilizer and soil conditioner.

Nutrient retention and higher nutrient availability were found in the soil after biochar addition, associated with higher CEC, surface acreage, and direct nutrient addition (Glaser *et al.*, 2002). In line with opinions, Murtaza *et al.* (2021) suggested that biochar is a recalcitrant carbon product of biomass produced through the pyrolysis process. The recalcitrant properties of biochar are distinctive and important to the long-term improvement of soil properties through the effects of their residues that can persist for a long time in the soil. Beusch (2021), biochar can contribute to soil fertility and crop productivity, and Kalu *et al.* (2021) increase water availability and lower soil BD.

Applying biochar can improve soil properties and long-term crop productivity compared to other organic amendments. In other organic amendments, research results by Suri and Yudono (2020) showed that giving *Cromolaena odorata* as compost can increase plants' nutrition and nutrient uptake. Agustina *et al.* (2019) argued that using *Chromolaena odorata* and rice straw can maintain soil quality. Mahmoud *et al.* (2009) also showed rice straw compost could improve soil quality. The same was discovered by Gewaily (2019), that straw compost can increase potassium availability.

Furthermore, the results of the study by Yin *et al.* (2022) showed the weight of the soil contents decreased, and the water-holding capacity and total porosity increased in the soil with rising rice husk ash. Adding rice husk ash neutralizes soil acidity and increases plant nutrient availability. Research by Lestari and Rachmawati (2020) showed that applying rice husk ash (4 t/ha) can increase rice growth and reduce salinity stress.

Previously, several studies on the effect of biochar singularly have been carried out, including on the P-availability (Zhang *et al.*, 2022), improvement of soil physical properties (Aslam *et al.*, 2014), chemical properties of the soil (Bista *et al.*, 2019), and some soil characteristics (Yadav *et al.*, 2018), soil aggregate stability (Ebido *et al.*, 2021), and to the development of rice roots (Kartika *et al.*, 2021). However, no research has been found on

how the role of biochar was given together with other organic amendments in acid sulphate soil. Therefore, it is necessary to study how the effect of biochar is given together with organic amendments to *Chromolaena odorata* and rice straw. This research is expected to contribute to improving the nature of acid sulphate soil in West Kalimantan, which has been one of the types of soil widely used as an agricultural activity for rice crops. In connection with this background and literature review above, this study aimed to know the role of rice husk biochar combined with amendments to *Chromolaena odorata*, rice straw, and rice husk ash in improving the properties of acid sulphate soil.

MATERIALS AND METHODS

Study Area

This research was conducted from September to November, 2021. The research site was carried out at the Laboratory of the Faculty of Agriculture, Universitas Panca Bhakti, Pontianak, West Kalimantan Province, Indonesia. The height of the study site is one m from above sea level, with an average temperature and humidity of 27.6 °C and 82.8%, respectively. The research position is located at a latitude of 2°05' N–3° 05' S and longitude of 108°30' E–144°10' W.

Experimental Design

The study was conducted in a greenhouse and arranged in a complete randomized design (CRD) with three replications. Biochar amendments consisted of nine kinds, namely: without amendments (control), *Chromolaena odorata* (10 t/ha), rice straw (10 t/ha), rice husk biochar (10 t/ha), rice husk ash (10 t/ha), *Chromolaena odorata* (10 t/ha) + rice husk biochar (5 t/ha), rice straw (10 t/ha) + rice husk biochar (5 t/ha), rice straw (10 t/ha) + rice husk biochar (5 t/ha). In all, 27 polybags are needed.

Research Procedures

The material in this study used acid sulphate soil taken on the experimental land of the West Kalimantan, Sungai Kakap, Agricultural Technology Assessment Center, and land farmers' work on rice crops. This area included the development of swampland agriculture which has built irrigation canals since 1980 and was currently designated as an Integrated Farmer Business Area. Soil picking was carried out randomly at a soil depth of 0-20 cm, then mixed, air-dried, and cleaned. Then put in polybags was as much as 8 kg.

Biochar was made by taking rice husk from grain harvested from rice plants grown directly in the acid sulphate soil of the Kakap River, West Kalimantan, Indonesia. First, the rice mill was taken the grain to obtain rice husk. Furthermore, the rice husk was placed on the pyrolysis device to be processed into biochar by incomplete combustion (limited air) in a reactor made of Pertamina drums.

Organic amendments to *Chromolaena odorata* were taken from around the field, then chopped to a size of 1-2 cm. Rice straw was taken from waste harvested rice grown and grown by farmers in acid sulphate soil and chopped within 1–2 cm. Rice husks were taken from waste from rice milling, then used as ash through burning in open areas to produce grayish-white ash.

Prepared acid sulphate soil introduced into 27 polybags until it reached 8 kg. Further, the dose of soil amendment was calculated based on the soil weight in the polybag area, mixed with the soil in each polybag to a soil depth of 20 cm, after which it was incubated with moisture content until it was closed to the field capacity for 30 days.

At 30 days after incubation, from each study unit, soil sampling of about 50 g was undisturbed to determine the weight of the soil moisture content accident. In addition, a disturbed soil sample of 100 g was taken from each research unit for analysis of chemical properties.

Observation parameters

The soil bulk density (BD) was determined by the clod method described by Blake & Harke. The total pore was calculated from the soil moisture content (v/v) at the matrix potential of 0 kPa. The available groundwater was estimated by reducing the soil moisture content in the matrix potential by -33 kPa (field capacity) with the soil moisture content in the matrix potential, which was -15 Mpa (wilting point). The soil moisture content at this potential matrix was determined using a pressure plate device. The strength of the soil was measured with a hand penetrometer (Daikie) at a soil depth of 15 cm.

The pH level was measured in a soil solution at a ratio of 1:2.5 (with deionized water), using a pH meter (Jenway 3305). Walkley and Black's wet oxidation method was used to determining organic C levels (Soil Survey Laboratory Staff). Total N levels were measured by the Kjeldhal (Bremner and Mulvaney) method. $Al_3 + and Fe_2 + were$ extracted with 1 M KCl (Barnhisel and Bertsch). CEC was extracted with 1 M NH₄Oac (buffer at pH 7.0), and the concentration of alkaline cations was measured using AAS (Shimadzu), P-available with Bray I.

Statistical Analysis

Statistical analysis was carried out on each observation parameter data to determine the effect of the amendments on the nature of acid sulphate soil. The data obtained were analyzed by analysis of variance (ANOVA) at 5% significant level. In addition, the test of least significant difference (LSD) at 5% significant level was used to determine the average value between treatments.

RESULTS AND DISCUSSION

Effect of various soil amendments on some physical soil properties

Applying biochar improved the soil's physical properties (Aslam *et al.*, 2014). Applying biochar and organic amendments influenced the changes in some of the observed soil physical properties. The statistical test is shown in Table 1. The soil BD was lower on all amendments granted. For example, in soils fed a single rice husk biochar (10 t/ha), soil BD decreased from 1.27 mg/m³ in the control soil to 1.15 mg/cm^3 . However, not significantly different from soils was given straw (10 t/ha) and a combination of straw + biochar, *Chromolaena odorata* + rice husk ash, and straw + rice husk ash. Furthermore, in the combination of *Cromolaena odorata* (10 t/ha) + rice husk biochar (5 t/ha), a BD of 1.13 mg/m^3 was obtained and was the lowest soil BD. Still, it was not significantly different from the soil, given a combination of straw (10 t/ha) + rice husk biochar 5 (t/ha) produced soil BD of 1.14 mg/m^3 . These results indicate an improvement in the physical soil properties if biochar was given together with organic amendments to *Chromolaena odorata*.

Applying various biochar amendments and their combinations increased the soil's total pore. The statistical test results are shown in Table 1. There was a significant difference in the total pore of the soil between the various amendments. Control soils had the lowest total pore (44.43%), contrasting with soils that provided various amendments. On soils with rice husk biochar (10 t/ha), the total soil pore increased to 54.21%, significantly different from other single amendments. Furthermore, a combination of *Cromolaena odorata* (10 t/ha) + rice husk biochar (5 t/ha) caused the total pore to increase to 56.73% and was the highest total pore.

The application of various amendments, both single and combined, affected the decrease in soil strength as measured as soil penetration resistance (N/cm²) at conditions of pF 0, pF 2, and pF 2.5, where the higher the soil matric suction (pF), the higher the soil penetration resistance. The that under saturated conditions (pF 0), the various soil amendments do not offer a noticeable difference. In pF 2 showed that the application of rice husk biochar (10 t/ha) can reduce soil penetration resistance from 310.00 N/cm² (control soil) to 223.33 N/cm², but not significant difference between *Chromolaena odorata* (10 t/ha) + rice straw or a combination of straw (10 t/ha) + rice husk biochar (5 t/ha). Furthermore, soils given a combination of *Cromolaena odorata* (10 t/ha) + biochar (5 t/ha) obtained the lowest penetration resistance (220.00 N/cm²), and this treatment was not significantly different from soils have given *Chromolaena odorata* (10 t/ha) + rice husk biochar (10 t/ha), rice husk biochar (10 t/ha), and a combination of rice straw (10 t/ha) + rice husk biochar (5 t/ha).

Similarly, in pF 2.5, the application of rice husk biochar (10 t/ha) could reduce the value of soil penetration resistance from 500.00 N/cm² (control soil) to 390.00 N/cm², and this treatment not significantly different from soils have given *Chromolaena odorata* (10 t/ha), rice straw (10 t/ha), rice husk ash (10 t/ha), and a combination of rice straw (10 t/ha) + rice husk ash (5 t/ha). Furthermore, the combination of *Cromolaena odorata* (10 t/ha) + rice husk biochar (5 t/ha) achieved the lowest

penetration resistance (340.00 N/cm²). Also, it was not significantly different from the combination of rice straw (10 t/ha) + biochar (5 t/ha) (360.00 N/cm²). It indicated that rice husk biochar had the potential to control soil strength. In line with Das *et al.* (2021), applying biochar into the soil could reduce soil strength.

Effect of rice husk biochar on some soil chemical properties

The application of several soil amendments and their combinations could influence some chemical properties of acid sulphate soil. The results of statistical tests in Table 2 showed that soil reaction (pH) of the soil increased in all treatments given compared to controls. For example, in the application of rice husk biochar showed an increase in pH from 3.36 (control soil) to 4.40 on soils given biochar (10 t/ha) but did not significantly different from other single amendments. Furthermore, the combined application of *Cromolaena odorata* (10 t/ha) + biochar (5 t/ha) obtained the highest increase in soil pH (4.58), in marked contrast to soils given rice husk ash but not significantly different from single amendments or other combinations.

Applying single and combined soil amendments affected to increase in soil C-organics. The statistical test results in Table 2 showed that the control soil had the lowest C-organic (1.94%), in stark contrast to other amended soils, except that the soils have given rice husk ash (10 t/ha) was not significantly different. The soil given a single amendment of rice straw (10 t/ha) obtained 4.58% C-organic, in stark contrast to the control soil and the soil given rice husk ash, but not significantly different from other soil amendments. Similarly, soils given a combination of *Chromolaena odorata* (10 t/ha) + biochar husk (5 t/ha) achieved the highest soil C-organic (4.93%) in marked contrast to control soils and soils fed with rice husk ash but did not significantly different from other soil amendments.

Applying various amendments and combinations to acid sulphate soil generally affected the increase in P-available. The results of statistical tests (Table 2) showed that control soils had the lowest P-available content (0.21%), significantly different from soils given various amendments. In soils fed with a rice husk biochar (10 t/ha), the P-available increased to 0.32%. Still, it was not significantly different from other combined or single-amended soils, except with soils that were given rice husk ash with a lower P-available (0.27%). The highest P-available was obtained on soils given a combination of *Chromolaena odorata* (10 t/ha) + biochar (5 t/ha) (0.34%), in stark contrast to control soils and those topped with rice husk ash but not significantly different from single amendments or other combinations.

The decrease in Al-dd and Fe-dissolve in acid sulphate soil also occurs due to applying various soil amendments and their combinations. Statistical tests (Table 2) showed that the highest Al-dd was obtained on control soil (3.84%), further decreasing and differing markedly from soils given various single and combined amendments. On single-amendment soils of rice husk biochar (10 t/ha),

Al-dd decreased to 3.16%, significantly different from other single-amended soils. Furthermore, the soil given the combination of *Chromolaena odorata* (10 t/ha) + rice husk biochar (5 t/ha) obtained the lowest Al-dd content (3.02%). It was not significantly different from the soil given a combination of rice straw (10 t/ha) + rice husk biochar (5 t/ha), but it was significantly different from other amended soils. In soluble Fe content, the application of rice husk biochar (10 t/ha) could reduce soluble Fe from 3.61% on control soil to 3.10% and was significantly different from other single amendments. And when the soil was given a combination of rice straw (10 t/ha) + rice husk biochar (5 t/ha). Still, it was not significantly different from soil that was given a combination of *Chromolaena odorata* (10 t/ha) + rice husk biochar (5 t/ha).

The CEC of land has also increased due to the application of various land amendments and their combinations. The statistical test is shown in Table 2 show that the control soil had the lowest CEC (6.64 Cmol/kg), not significantly different from soils fed with *Chromolaena odorata* and rice straw. On the other hand, on soils provided with rice husk biochar (10 t/ha), the soil CEC increased to 8.03 Cmol/kg, but it was not significantly different from soil given rice straw and rice husk ash.

In line with the results that have been put forward, Yamato *et al.* (2006) found that the application of biochar from *Acacia mangium* bark into the soil in Sumatra led to changes in soil chemical properties through increased pH, N-total and P-available, CEC, and lowered interchangeable Al. Liang *et al.* (2006) found that two mechanisms could create a larger CEC. First, a higher charged density per unit of surface acreage means a higher degree of oxidation of soil organic matter. Second, the presence of a higher surface area for cation absorption or the combined effect of both. Glaser *et al.* (2002) posit that oxidation of aromatic C and carboxyl formation are the main reasons for the high CEC. Liang *et al.* (2006) also found that the CEC per unit C was higher, and the load density was higher in carbon-black-rich Anthrosol compared to soils that were poor in carbon black. In addition, Anthrosol showed a higher surface acreage due to its higher carbon black concentration.

From the results, the great potential of rice husk biochar against improving soil properties increased when given together with the biomass *Chromolaena odorata*. It could happen because *Chromolaena odorata* contains higher humate and fulvic organic acids, so that it could encourage the complex mechanisms of organo-mineral organs and soil aggregation. In addition, *Chromolaena odorata* can highly suppress Al and Fe, thereby increasing the P availability in the soil due to the presence of organic acids released during decomposition. Research results by Suri and Yudono (2020) also found that using *Chromolaena odorata* compost could improve soil quality and nutrient uptake of lettuce plants.

Chromolaena odorata can contribute to the care of soil C-organic and physical properties due to a reasonably high polyphenol content that could inhibit the decomposition and mineralization of N organic matter, even though it had a low C/N ratio.

CONCLUSIONS

In conclusion, the amendment of rice husk biochar combined with amendments to *Chromolaena odorata* and rice straw improved soil physical properties, namely total soil pores, reducing soil content weight and strength. Its impact on soil chemical properties was that it could increase pH, C-organic, P-available, CEC, lower Al-dd, and Fe soluble. Therefore, applying biochar and organic amendments to *Chromolaena odorata* and rice straw could improve the properties of acid sulphate soil. Consequently, we recommend increasing the carrying capacity of acid sulphate soil against plant growth. It can be done by enhancing soil properties through the organic amendments application of *Chromolaena odorata* or straw enriched with rice husk biochar.

ACKNOWLEDGEMENTS

We thank the Institute for Research and Community Service, Universitas Panca Bhakti, which has given support for research funds. In addition, we thank the Faculty of Agriculture, Universitas Panca Bhakti, which has provided loans for facilities in the form of equipment for research.

REFERENCES

- Agustina, R., Jumadi, R., Firmani, U. and Abdul Faisal, R. H. (2019). Effects of decomposition rate of *Chromolaena odorata* and straw rice in fresh and compost form to the growth and yield of rice. *IOP Conf Ser Earth Environ Sci.* **250**: 1–5.
- Aslam, Z., Khalid, M. and Aon, M. (2014). Impact of biochar on soil physical properties. *Sch J Agric Sci.* **4**: 280–4.
- Beusch, C. (2021). Biochar as a soil ameliorant: How biochar properties benefit soil fertility A review. J Geosci Environ Prot. 9: 28–46.
- Bista, P., Ghimire, R., Machado, S. and Pritchett, L. (2019). Biochar effects on soil properties and wheat biomass vary with fertility management. *Agronomy*. **9**: 1-10.
- Das, S. K. and Das, S. K. (2015). Acid sulphate soil: Management strategy for soil health and productivity. *Pop. Kheti.* **3**: 3–8.
- Das, S., Mohanty, S., Sahu, G., Rana, M. and Pilli, K. (2021). Biochar: A sustainable approach for improving soil health and environment. In: Soil Erosion - Current Challenges and Future Perspectives in a Changing World. pp. 1-18.
- Ebido, N. E., Edeh, I. G., Unagwu, B. O., Nnadi, A. L., Ozongwu, O. V., Obalum, S. E. and Igwe, C.
 A. (2021). Rice-husk biochar effects on organic carbon, aggregate stability and nitrogen-fertility of coarse-textured Ultisols evaluated using *Celosia argentea* growth. *Sains Tanah.* 18: 177–87.
- Gewaily, E. (2019). Impact of compost rice straw and rice straw as organic fertilizer with potassium treatments on yield and some grain quality of Giza 179 rice variety. *J Plant Prod.* **10**: 143–51.
- Glaser, B., Lehmann, J. and Zech, W. (2002). Ameliorating physical and chemical properties of highly weathered soils in the tropics with charcoal A review. *Biol Fertil Soils*. **35**: 219–30.

- Kalu, S., Simojoki, A., Karhu, K. and Tammeorg, P. (2021). Long-term effects of softwood biochar on soil physical properties, greenhouse gas emissions and crop nutrient uptake in two contrasting boreal soils. *Agric Ecosyst Environ.* **316**: 107454.
- Kartika, K., Sakagami, J. I., Lakitan, B., Yabuta, S., Akagi, I., Widuri, L. I., Siaga, E., Iwanaga, H. and Nurrahma, A. H. I. (2021). Rice husk biochar effects on improving soil properties and root development in rice (*Oryza glaberrima* Steud.) exposed to drought stress during early reproductive stage. *AIMS Agric Food.* 6: 737–51.
- Kocsis, T., Ringer, M. and Biró, B. (2022). Characteristics and applications of biochar in soil–plant systems: A short review of benefits and potential drawbacks. *Appl Sci.* **12**: 1–16.
- Lestari, M. F. and Rachmawati, D. (2020). The effect of rice husk ash on growth of rice (*Oryza sativa* L. 'Sembada Merah') on salinity stress. *AIP Conf Proc.* **2260**: 1–6.
- Liang, B., Lehmann, J., Solomon, D., Kinyangi, J., Grossman, J., O'Neill, B., Skjemstad, J. O., Thies, J., Luizão, F. J., Petersen, J. and Neves, E. G. (2006). Black carbon increases cation exchange capacity in soils. *Soil Sci Soc Am J.* **70**: 1719–30.
- Mahmoud, E., Ibrahim, M., Robin, P., Akkal-Corfini, N. and El-Saka, M. (2009). Rice straw composting and its effect on soil properties. *Compost Sci Util.* **17**: 146–50.
- Murtaza, G., Ahmed, Z., Usman, M., Tariq, W., Ullah, Z., Shareef, M., Iqbal, H., Waqas, M., Tariq, A., Wu, Y., Zhang, Z. and Ditta, A. (2021). Biochar induced modifications in soil properties and its impacts on crop growth and production. *J Plant Nutr.* 44: 1677–91.
- Shamshuddin, J., Elisa Azura, A., Shazana, M. A. R. S., Fauziah, C. I., Panhwar, Q. A. and Naher, U. A. (2014). Properties and management of acid sulfate soils in Southeast Asia for sustainable cultivation of rice, oil palm, and cocoa. *Advances in Agronomy*. **124**: 91-142
- Sovova, S., Enev, V., Smilek, J., Kubikova, L., Trudicova, M., Hajzler, J. and Kalina, M. (2021). The effect of biochar application on soil properties and growth of the model plant *Zea mays*. *Ecocycles*. **7**: 46–54.
- Suri, A. M. and Yudono, P. (2020). Effects of *Chromolaena odorata* compost on soil and nutrient uptake of lettuce (*Lactuca sativa*). *Planta Trop J Agro Sci.* **8**: 33–8.
- Yadav, N. K., Kumar, V., Sharma, K. R., Choudhary, R. S., Butter, T. S., Singh, G., Kumar, M. and Kumar, R. (2018). Biochar and their impacts on soil properties and crop productivity: A review. *J Pharmacogn Phytochem Harpole*. 7: 49–54.
- Yamato, M., Okimori, Y., Wibowo, I. F., Anshori, S. and Ogawa, M. (2006). Effects of the application of charred bark of *Acacia mangium* on the yield of maize, cowpea and peanut, and soil chemical properties in South Sumatra, Indonesia. *Soil Sci Plant Nutr.* 52: 489–95.
- Yin, M., Li, X., Liu, Q. and Tang, F. (2022). Rice husk ash addition to acid red soil improves the soil property and cotton seedling growth. *Sci Rep.* 12: 1–9.

Zhang, Y., Chen, H., Xiang, J., Xiong, J., Wang, Y., Wang, Z. and Zhang, Y. (2022). Effect of ricestraw biochar application on the acquisition of rhizosphere phosphorus in acidified paddy Soil. *Agronomy*. **12**: 1-13.

Table 1. Effect of rice husk biochar with soil amendments on some physical properties of acid sulphate soil at one month after incubation.

Collomon durante	BD	Total	Penetration resistance (N/cm ²)				
Soll amendments	(mg/m^3) pores (%)		pF 0	pF 2	pF 2.5		
Control	1.24 e	44.43 a	36.67 b	310.00 e	500.00 d		
C. odorata (10 t/ha)	1.16 c	52.17 e	20.00 a	230.00 abc	403.33 bc		
Rice straw (10 t/ha)	1.17 bc	53.27 f	16.67 a	243.33 bc	393.33 b		
Rice husk biochar (10 t/ha)	1.15 b	54.21 g	14.33 a	223.33 ab	390.00 b		
Rice husk ash (10 t/ha)	1.19 d	47.30 b	20.00 a	270.00 d	403.33 bc		
C. odorata (10 t/ha) +	1.13 a	56.73 i	10.00 a	220.00 a	340.00 a		
biochar (5 t/ha)							

Rice straw (10 t/ha) + rice	1.14 ab	55.57 h	13.33 a	223.00 ab	360.00 a
husk biochar (5 t/ha)					
C. odorata (10 t/ha) + rice	1.15 b	48.90 c	16.67 a	240.00 bc	420.00 c
husk ash (5 t/ha)					
Rice straw (10 t/ha) + rice	1.16 bc	50.30 d	20.00 a	246.67 с	403.33 bc
husk ash (5 t/ha)					

Remarks: The average numbers followed by the same letter in the same column show no significant difference in the LSD test at 5% significant level.

Table 2. Effect of rice husk biochar with soil amendments on some chemical properties of acid sulphate soil at one month after incubation.

		C-	Р-	Al-dd	Soluble	CEC
Soil amendments	pH H ₂ O	organic	available	(me/100	Fe	(me/100
		(%)	(%)	g)	(%)	g)
Control	3.36 a	1.94 a	0.21 a	3.84 e	3.61 d	6.64 a
C. odorata (10 t/ha)	4.06 bc	4.22 b	0.29 bc	3.31 c	3.28 c	7.15 ab
Rice straw (10 t/ha)	4.28 bc	4.58 b	0.30 bc	3.42 cd	3.34 cd	7.32 abc
Rice husk biochar (10 t/ha)	4.40 bc	4.09 b	0.32 cd	3.16 b	3.10 abc	8.03 c
Rice husk ash (10 t/ha)	3.98 b	2.78 a	0.28 b	3.51 d	3.34 cd	7.76 bc
C. odorata (10 t/ha) + rice	4.58 c	4.93 b	0.34 d	3.02 a	2.94 ab	10.04 e
<mark>husk</mark> biochar (5 t/ha)						
Rice straw (10 t/ha) + rice	4.48 bc	4.75 b	0.33 d	3.08 ab	2.91 a	9.67 de
husk biochar (5 t/ha)						
C. odorata (10 t/ha) + rice	4.26 bc	4.70 b	0.31 cd	3.37 c	3.25 c	9.17 d
husk ash (5 t/ha)						
Rice straw (10 t/ha) + rice	4.26 bc	4.73 b	0.31 cd	3.47 d	3.22 bc	9.20 d
husk ash (5 t/ha)						

Remarks: The average numbers followed by the same letter in the same column show no significant difference in the LSD test at 5% significant level.

Manuscipt acceptance : 29 Oktober 2022





Pembayaran : 31 Oktober 2022

_		~ "									9	
=	M	Gmail		Q	isent	×	12	Aktif • ⑦ 😝		(C)	UPI	3 (
990 ail	1	Tulis		÷				8	dari 24	٢	>	
		Kotak Masuk	5.340	۲	Agusalim Masulili «agusalim@upb.ac.id»			@ 31 Okt 2022 14.01	☆	ر)	÷	
ŧt	☆	Berbintang			Dear.							
8	0	Ditunda			Attached herewith is proof of payment for our paper "Role of biochar amendmer	nts in improving the prop	erties of acid sul	phate soil - Agusalim Masulili, Sutikarini, F	≀ini Sury	yani, Ida	Ayu	
es	>	Terkirim			Suci, ismail Astar, Harol Dominikus Bancin and Palman							
	۵	Draf	4		Thank you, Best regards.							
rt	~	Selengkapnya			Agusalim Masulili							
					Panca Bhakti University, Pontianak, Indonesia							
	Lab	el	+									
					Satu lampiran • Dipindai dengan Gmail 🛈						@ +	
					And the address and the Address Brown and the Address Address Address Address Address Address Address Address A							
					The second							
					Transfer_Agusali							
				G	Gaurav Publications <researchoncrops@gmail.com></researchoncrops@gmail.com>			31 Okt 2022 14.16	☆	4	:	
								Nonaktifkar	n untuk:	Ingaris		
					A mggna - y mooneara - terjentatikan pesan				1000000		0	

Bukti pembayaran:

M Acceptance of paper for publicat X S transfer_Agu	usalimPaper (1).pdf X 🛇 Invoice 11147 (1).pdf X 🛇 AL 11147 (1).pdf X +	✓ - □ ×
\leftarrow \rightarrow C O File C:/Users/asus/Downloads/t	transfer_AgusalimPaper%20(1).pdf	ie 🕁 🔒 📴 🕒 🌲 🛙 🦺 :
	1 / 1 - 100% + 🗄 👌	± ē :
	<section-header><form><form></form></form></section-header>	
\bigcirc Type here to search	H 🗷 🗊 🗛 💽 🔚 💽 🖬 🚺	🌰 30°C Berawan へ ট 句) ENG 11:09 AM 🖓

Pemberitahuan penerimaan pembayaran biaya publikasi: 31 Oktober 2022

Accep	tance of paper for publicati 🗙	S transfer_Agusalimi	laper (1).pdf 🛛 🗙 📔 😵 Invoice 11147 (1).pdf 💦 🗙 📔 🏵 AL 11147 (1).pdf	× +	~ - 🗆 ×
$\leftarrow \rightarrow$	C a mail.google.com/m	nail/u/3/?ogbl#sent/	FMfcgzGqRPvCwFrMhQtzWbzpJVnztflX		🖻 🖈 🦰 🦝 🖷 🧶 i
=	M Gmail	Q	in:sent	× 辈	• Aktif - 🔿 🏶 🏭 🎯 UPB 🍥
Mail	/ Tulis	~			8 dari 24 < >
Chat	G Kotak Masuk ☆ Berbintang	5.340	🔛 transfer_Agusali 🥢		•
:8:	O Ditunda	G	Gaurav Publications <researchoncrops@gmail.com></researchoncrops@gmail.com>		31 Okt 2022 14.16 🚖 🕤 🚺
spaces	> Terkirim				Nooskifkan untuk longris
0	Draf	4	XA inggris + > indonesia + ierjemankan pesan		
Middt	 Selengkapnya 		Dear Agusalim,		
	Label	+	Received the copy of the payment document. You will be informed soon after the receipt of payment in our bank account I hank you. Dr. Vedgal Singh Chief Editor see Noted with thanks. Thank you for your response. Thank you for the Grant Balas of Teruskan	ne information.	
					11:14 AM
4	Type here to search		- H: 🐼 😐 🚧 💽 🧰 🚺		🌰 31°C Berawan 🥎 💮 🗇 ENG 12/22/2022

Galley copy of paper for publication (10.11147) : 8 Desember 2022

fourub	e 🛃 Maps																
=	M Gmail		Q 1	elusuri dalam email						井	• Aktif 👻	0	19 3			UP	в
22-	Tulis			₽ () Ĥ		G D	D :						1–50 dar	i 5.505	<	>	
ii 1	- Kotak Masuk	5.340		VEDPAL, saya 3	Galle	y copy of pape	er for public	cation (10. 11147	') - Thanks for y	our confirmation	to consider galley co	py for fin	nal publi	c		8 Des	
at	A Berbintang					10. 11147 Galle	ey										l
a es	Ditunda			Jurnal Ilmu Pertani.	[JIPI]	New notificati	tion from Jur	rnal Ilmu Pertar	nian Indonesia -	You have a new	notification from Jurn	al Ilmu P	ertania	n I		7 Des	
4	Draf	4		Grommarky logishts	[JIP]	weekky etete	aala autaad	dod: 50% Off	Your writing of	to from loot woo	k are amazing! Cramp		VEMPE	0		6 Des	
et	 Selengkapnya 			Grammany insights	tour	weekly stats +	- sale extend	ded: 50% Off! -	tour writing sta	ats from last wee	k are amazing! Gramn	harly NO	VEMBE	R		o Des	
	l - h - l			Google	Notifi	kasi keamanar	n - Login ba	aru di Windows	agusalim@upb.	ac.id Kami mene	mukan adanya login b	aru ke A	kun Go	0		5 Des	
	Label	+		Dr. Iman Rusmana	[JIPI]	Password Res	set - Your pa	assword has be	en successfully	reset for use wi	th the Scientific Journ	als of IP	B Unive	rs		5 Des	
				Dr. Iman Rusmana 2	[JIPI]	Password Res	set Confirma	ation - We have	e received a req	uest to reset you	ir password for the Sc	ientific J	lournals	0		5 Des	
				Jurnal Ilmu Pertani.	[JIPI]	New notificati	ion from Jur	rnal Ilmu Pertar	nian Indonesia -	You have a new	notification from Jurn	al Ilmu P	ertania	n I		5 Des	
				Asian Institute of .	Dece	mber Issue - E	Education Re	esearch Publica	ation Invitation -	- ISSN 2621-5799	Call for Papers: Vol.5	No.4 (De	ecembe	r		4 Des	
				Scholastica	[The	Abstract] Auth	hor support	insights + new	publishing anal	ytics - In this iss	ue: Authorship dos an	d don'ts,	, a guide	ə	31	0 Nov	
				Grammarly Insights	Just	for you: This	week's stat	ts + your Cybe	er Monday offe	r - Let's get bacl	k to it! Make sure you'	re logge	d in. Gra)	29	Nov	
				Dion Septian (Class.	. Terla	mbat Dikirim	ıkan: "Tuga:	s tulis resume	materi 5, kum	pulkan" - Hai A	Agusalim, Dion Septiar	n baru sa	ija meng	g	21	7 Nov	
				Adi Saputra - 2989 .	3 Terla	mbat Dikirim	ıkan: "Tugas	s tulis tangan	materi 5, kump	pulkan" - Hai A	gusalim, Adi Saputra	- 2989 bi	aru saja		23	3 Nov	
О т Salley	Type here to search	+			× •						▲ 31°C	Berawan		<u>⊡</u> ⊄»)	ENG 12	2:06 PM 2/22/20	
O T Galley → ouTub	Type here to search copy of paper for publicat: X C (a) mail.google.com, re (C) Maps	+ a/mail/u/0/?ta	ib=rm&og	H Z 🗊	pGkfTCSTCsZXgc	knkSsNr					▲ 31*C	Berawan		∰ (1)) ⊛	ENG 12	2:06 PM 2/22/202	2
Ω T Galley →	Copy of paper for publicant X C end a mail.google.com, Re R Maps M Gmail	+ /mail/u/0/?ta	ib=rm&og	다. 조 🗊	pGkITCSTCsZXge	knkSsNr				T	● 31*C ● Aktif ▼	@	• ^ (₫ (1)) ₫ ₫ ∅ ∅ 	ENG 12	2:06 PM 1/22/202	2
Colley → touTub	Type here to search copy of paper for publican × C mail.google.com, Maps G Gmail Tulis	+ /mail/u/0/?ta	b=rm&og Q T ¢	Eli aluminbox/EMfcgzGrbRX; Felusuri dalam email	pGkITCSTCsZXgc	knkSsNr				#	● 31°C	@	 33 dar 	 ₫ (1-3) 	ENG 12 ✓	2:06 PM (722/202	
Q T Galley → ouTub	Type here to search copy of paper for publican × C mail.google.com, Maps G Gmail Tulis Kotak Masuk	+ //mail/u/0/?ta	b=rm&og Q T +	Eliterinbox/EMIcgzGrbRX;	pocktTCSTCsZXge	knkSsNr @	 Image: A state of the state of	11147)	tternal) Kotak M	子 fasuk ×	● 31°C	Berawan	 (2) (2) (2) (3) (4) 	 ■ (1-0) ● (1-0)	ENG 12	2:06 PM (22/202	2
Galley → buTub	Type here to search	+ +/mail/u/0/?ta	b=rm&og Q T ¢	Eli al Elistrico del ante en elistrico del ante elistrico del ante en elistrico del ante	GektTCSTCsZXge ⇒GektTCSTCsZXge f paper for ditor@gmail.com>	knkSsNr @ publicati	₽ i	11147)	iternal) Kotak M	T#	Aktif -	Rerawan	 33 dar es 19.09 	 (a) (b) (c) <li(c)< li=""> <li(c)< li=""> <li(c)< li=""> (c)</li(c)<></li(c)<></li(c)<>	ENG 12	2:06 PM (22/202 UP > 2 2 3 2 2 3 2 3 3 2 3 3 3 3 3 3 3 3 3	1
C T Galley → ouTub	Tulis Kotak Masuk Berbintang O Ditunda	+ + 5.341	b=rm&og Q T ¢	Elistinbox/EMIcgxGrbRX; Felusuri dalam email Calley copy of VEDPAL SINGH -rocec kepada saya =	pGktTCSTCsZXge	knkSsNr & D	₽ :	11147)	aternal) Kotak M	子 fasuk ×	Aktif -	Kam, 8 D	33 dar	 ● (1)) ● ●<		206 PM /22/202 UP	
Galley → t t	Tulis Kotak Masuk Kotak Masuk Ditunda Terkirim Di Do f	+ + 5.341	b=rm&og Q T ¢	Image: Second	constant of the second se	knkSsNr 2 D 2 publicati	₽ i	11147)	sternal) Kotak M	王上 tasuk ×	Aktif -	Contraction (Contraction (Contraction)) (Contraction) (Con	33 dar es 19.09	● tv)) ● tv) i 5.506 ☆	ENG 12	2:06 PM /722/202 UP 2: 2: 2: 2: 2:	1
Galley → ouTub	type here to search C C C C C C C C C C C C C C C C C C	+ //mail/u/0/?ta 5.341	b=rm&og Q T ÷	Image: Second	construction	knkSsNr ? publicati	 ion (10. 	11147)	H OL COARS IS	포 tasuk ×	Aktif -	Rerawan (2) Kam, 8 D Not	A (33 dar as 19.09	 (€) (€)	ENG 12	206 PM //22/202 ↓ ↓ ↓ ↓ ↓ ↓ ↓ ↓ ↓ ↓	1
Galley → fourtube t t t t	type here to search coopy of paper for publicar: × C a mail.google.com, a mail.google.com, b Maps C Tulis Kotak Masuk A Berbintang C Ditunda F Terkirim D Draf Selengkapnya Labad	+ //mail/u/0/?ta 5.341	ab=rm&og Q T E		O	knkSsNr knkSsNr Couperation	ion (10. eson eson	ed in RESEARC	Kotak M Kotak M :H ON CROPS Jo	子上 fasuk ×	 Aktif ~ Aktif ~ 4 (December) 2022. 	(2) Kam, 8 D Nor	 A (33 dar 33 dar 	* * i 5.506 \$ \$ \$	ENG 12 안 소 · · ·	206 PM //22/202 UP > : : :	
Calley → fouTube fouTube fouTube fouTube fouTube fout fou	type here to search	+ //mail/u/0/?ta 5.341 4 +	ab=rm&og Q T ¢	In the corrected copy show the corrected copy show the corrected copy show the comp show the comp show the corrected copy show the copy show the corrected copy show the copy show the copy show the corrected copy show the	AGUETCSTC-32Xge G	knkSsNr knkSsNr Publicati Terjemahkan pe of your paper for i, if any The core or your paper and allo	ion (10. csan ccsan	ed in RESEAR with the heightight	Motik M Scitik M SH ON CROPS Jo ed.	국는 tasuk × aurnal Vol. 23, No.	Aktif	? Noi	 A (General Activity 33 dar 33 dar 33 dar 	 ● ●	ENG 12	206 PM //22/202 UP > 22 : :	
C Galley → fouTubb fouTubb fouTubb fouTubb fourtu	type here to search	+ //mail/u/0/?ta 5.341 4 +	b=m&og Q T ¢	I I I I I I I I I I I I I I I I I I I	control contro control control control control control co	knkSsNr knkSsNr compared to the second secon	ion (10. csan be publisher rrections sho achment with he article. The	ed in RESEARd uid be highlighte in three days. ereofore, read the	H ON CROPS jo rd	파고 파고 No. 23, No.	31*C Aktif ~ 4 (December) 2022. ns, if any.	② Kam, 8 D Note:	 A (33 dar 33 dar 33 dar 	 (1) (2) (2) (3) (4) (5) (6) (7) (7)	ENG 12	206 PM (722/02 UP) 2 2 2 2 2 2 2 2 2 2 2 2 2	
P T Galley → fouTubo iii ii	type here to search	+ //mail/u/0/?ta 5.341 4 +	b=m8og	It is a set of the se	constant of the result of	knkSsNr knkSsNr comparenter forjemahkan pe forjemah	ion (10. csan o be published rirectors should achiment with he article. The	ed in RESEARC uid be highlight in fitnee days. orefore, read the	Hender Kotak M Kotak M CH ON CROPS jo d	T 파고 (Sasuk ×) nurnal Vol. 23, No. refully for correctio	Aktif	⑦ ⑦ Nore	 • • • (33 dar 33 dar 	 ● (10) ● ●<	ENG 12	206 PM (722/02) □ □ □ □ ·	
Control of the set of	type here to search	+ //mail/u/0/?ta 5.341 4 +	■		constant of the result of	knkSsNr knkSsNr comparenter forjemahkan pe forjemah	ion (10. csan o be published arrectors show a christian with he article. The	ed in RESEARC uid be highlight in fitnee days.	Hender Kotak M Kotak M CH ON CROPS jo d	Tasuk × nurnal Vol. 23, No. refully for correctio	Aktif Aktif	Image: Second	233 dar 233 dar	 (1) (2) (2) (3) (4) (5) (6) (7) (7) (7) (8) (9) (9)	ENG 12	206 PM /22/202 ↓ ↓ ↓ ↓ ↓ ↓ ↓ ↓ ↓	
	Tulis Kotak Masuk Kotak Masuk Kotak Masuk Kotak Masuk Kotak Masuk Kotak Masuk Selenjkapnya Label	+ + 5.341 4 +	b=m&ga 1 ¢	Image: Section 2016 Image: Section 2016 Imag	Compared attor DOI	knkSsNr knkSsNr comparison compar	ion (10. csan o be publishe rrections sho actiment with he article. The	ed in RESEARC vid be highlighte in three days.	tterna) Ketak M CH ON CROPS Jo d.	Issuk × nurnal Vol. 23, No. refuilly for correctio	Aktif	(2) Kam, 8 D Note:	 A C C	 (a) (1) (a) (1) (a) (1) (b) (1) (c) (1)	ENG 12 企 全	206 PM (7227202 ↓ (7227202 ↓ (7227202 ↓ (7227202 ↓ (7227202 ↓ (7227202 ↓ (7227202 ↓ (7227202 ↓ (7227202 ↓ (7227202) ↓ (7227202) ↓ (7227202) ↓ (7227202) ↓ (7227202) ↓ (72270) ↓ (722702) ↓ (722702) ↓ (722702) ↓ (722702) ↓ (72270) ↓ (72270) ↓ (72270) ↓ (72270) ↓ (72270) ↓ (72270) ↓ (72270) ↓ (72270) ↓ (72270) ↓ (72270) ↓ (72270) ↓ (72270) ↓ (72270) ↓ (72270) ↓ (7270)	
C Galley → KouTub Hill 1 1 1 1 1 1 1 1 1 1 1 1 1	Tulis Kotak Masuk Kotak Masuk Kotak Masuk Kotak Masuk Selengkapnya Label	+ + 5.341 4 +	b=rm&og		Compared atter DOI	knkSsNr knkSsN	ion (10. esan o be publishe rrections sho achment with he article. The	ed in RESEARC vid be highlight in three days. orefore, read the	anema) Kotak M CH ON CROPS jo rd	szuk × tasuk × rurnal Vol. 23, No.	Aktif	(2) Kam, 8 D Note:	a A (33 dar 33 dar as 19.09	 (1) (2) (3) (4) (5) (6) (7) (7)	ENG 12 企 全	226 FW /22/202 ↓ ↓ ↓ ↓ ↓ ↓ ↓ ↓ ↓ ↓ ↓ ↓ ↓	
C Galley → fouTub iii iii iii iii iii iii iii i	Type here to search	+ (mail/u/0/?ta 5.341 4 +	b=rm8tog		Construction C	knkSsNr knkSsNr knkSsNr knkSsNr publicati ferjemahkan pe of your paper to f, il any The corough email all and mail ①	ion (10. esan o be publishe rections sho he article. The	ed in RESEARC wid be highlighten orefore, read the	Remar Kotak M CH ON CROPS jo d.	Tasuk ×	Aktif	The second se	A (33 dar as 19.09	 ● (1)) ● ●<	ENG 12	226 FM /22/202 ↓ /22/20 ↓ /22/20 /22/20 ↓ /22/20 /22/20 /2 / /22/20 / / / / / / / / / / / / /	

Improving the properties of acid sulphate soil

Res. Crop. **23** (4): (2022) *Printed in India*

Role of biochar amendments in improving the properties of acid sulphate soil

AGUSALIM MASULILI*, SUTIKARINI, RINI SURYANI, IDA AYU SUCI, ISMAIL ASTAR, HARDI DOMINIKUS BANCIN AND PAIMAN¹

Department of Agrotechnology, Faculty of Agriculture Panca Bhakti University, Pontianak 78113, West Kalimantan, Indonesia *(e-mail: <u>agusalim@upb.ac.id</u>) (Received: October 03, 2022 / Accepted: November 28, 2022)

ABSTRACT

Acid sulfate soils have potential for agricultural development. However, this soil has various properties problems that can inhibit plant growth. One way to improve soil properties is through the application of biochar amendments. The study aimed at knowing the role of rice husk biochar combined with amendments to Chromolaena odorata, rice straw, and rice husk ash to improve the properties of acid sulphate soil. The study was conducted in a greenhouse and arranged in a complete randomized design (CRD) with three replications. Biochar amendments consisted of nine kinds, namely, without amendments (control), C. odorata (10 t/ha), rice straw (10 t/ha), rice husk biochar (10 t/ha), rice husk ash (10 t/ha), C. odorata (10 t/ha)+rice husk biochar (5 t/ha), rice straw (10 t/ha)+rice husk biochar (5 t/ha), C. odorata (10 t/ha)+rice husk ash (5 t/ha), and rice straw (10 t/ha)+rice husk ash (5 t/ha). The results showed that the amendment of rice husk biochar combined with amendments to C. odorata and rice straw, respectively, had a good effect on improving soil physical properties, namely, total soil pores, reducing soil content weight and soil strength. Its effect on soil chemical properties was that it could increase pH, C-organic, P-available, cation exchange capacity (CEC), lower Al-dd and Fe soluble. The research findings showed that applying biochar and organic amendments to Chromolaena odorata and rice straw could potentially improve the

¹Department of Agrotechnology, Faculty of Agriculture, Universitas PGRI Yogyakarta, Yogyakarta 55182, Indonesia.

properties of acid sulphate soil. In future research, we recommend increasing the carrying capacity of acid sulphate soil against plant growth, and it can be done by improving soil properties through the organic amendments application of *C. odorata* or straw enriched with rice husk biochar.

Key words: Chromolaena odorata, rice husk ash, rice husk biochar, rice straw, soil properties

INTRODUCTION

Acidic sulphate soils are the common name for soils and sediments containing iron sulfide, namely, pyrite (Das and Das, 2015). In agricultural development, efforts to reclaim acid sulphate soils do not always result in soil productivity since improvements in soil properties after reclamation are not achieved. So far, efforts have been encouraged to overcome existing obstacles after reclamation, including inorganic fertilizers (urea, TSP and KCl) and liming (dolomite). However, rice crop productivity in this field has not been optimal.

Inorganic fertilizers increase over time and can even decrease land quality if inputs are high and intensive. Liming can only cope with short-term pH and is often done repeatedly. As a result, high costs are needed, which becomes a burden for farmers and will threaten the sustainability of land productivity. Therefore, it is necessary to manage land based on land amendments that can improve the nature of sustainable soils.

Reclamation can result in changes in soil properties and is related to the potential for pyrite oxidation, affecting the increase in soil packaging and the solubility of ions such as Al, Fe, Mn and sulphate. On the other hand, this soil has a high clay content which affects the increased weight of the contents. Therefore, the development of acid sulphate soil for agriculture after reclamation requires proper management of soil amendments to achieve the formation of steady cultivated land and sustainable farming. Das and Das (2015) suggested that acid sulphate soil having a high pH could inhibit plant growth. In addition, when this soil is drained, it can affect an increase in sulfuric acid and the release of iron, aluminum, and other heavy metals. Shamshuddin *et al.* (2014) stated that plants grown in acidic sulphate soils could not grow well due to low pH and high Al content.

Biochar is one of the soil amendment materials with the long-term potential to maintain soil fertility (Ogunleye *et al.*, 2019; Jaswal *et al.*, 2022). The results of experts' investigations into the phenomenon of terra preta show that soil fertility is stable in the Amazon basin due to the presence of black carbon, the result of the management of the Amerindian nation 500-2,500 years ago. This carbon black is nothing but biochar immersed in the soil. Murtaza *et al.* (2021) found that 4.0 t/ha cotton biochar improved physical, chemical, soil biological soil fertility properties and peanut productivity in Alfisol. Zhang *et al.* (2022) have proven that biochar provides effects such as liming and providing nutrients that lead to soil improvement. Kocsis *et al.* (2022) stated that biochar had a high porosity and could improve the physico-chemical and biological properties of the soil. In line, Sovova *et al.* (2021) found that biochar had a good influence on improving the soil properties of regosol. Further, Das *et al.* (2021) suggest the application of biochar to the soil can provide agronomic benefits, support crop yield improvement, and improve soil quality and health. In addition, biochar has been shown to act as a fertilizer and soil conditioner.

Nutrient retention and higher nutrient availability were found in the soil after biochar addition, associated with higher CEC, surface acreage and direct nutrient addition (Glaser *et al.*, 2002). In line with opinions, Murtaza *et al.* (2021) suggested that biochar was a recalcitrant carbon product of biomass produced through the pyrolysis process. The recalcitrant properties of biochar are distinctive and important to the long-term improvement of soil properties through the effects of their residues that can persist for a long time in the soil. Beusch (2021), biochar can contribute to soil fertility and crop productivity, and Kalu *et al.* (2021) increase water availability and lower soil BD.

Applying biochar can improve soil properties and long-term crop productivity compared to other organic amendments. In other organic amendments, research results by Suri and Yudono (2020) showed that giving *C. odorata* as compost can increase plants' nutrition and nutrient uptake. Agustina *et al.* (2019) argued that using *C. odorata* and rice straw could maintain soil quality. Mahmoud *et al.*

(2009) also showed rice straw compost could improve soil quality. The same was discovered by Gewaily (2019), that straw compost can increase potassium availability.

Furthermore, the results of the study by Yin *et al.* (2022) showed the weight of the soil contents decreased, and the water-holding capacity and total porosity increased in the soil with rising rice husk ash. Adding rice husk ash neutralizes soil acidity and increases plant nutrient availability. Research by Lestari and Rachmawati (2020) showed that applying rice husk ash (4 t/ha) could increase rice growth and reduce salinity stress.

Previously, several studies on the effect of biochar singularly have been carried out, including on the P-availability (Zhang *et al.*, 2022), improvement of soil physical properties (Aslam *et al.*, 2014; Yin *et al.*, 2018), chemical properties of the soil (Bista *et al.*, 2019), some soil characteristics (Yadav *et al.*, 2018), soil aggregate stability (Ebido *et al.*, 2021) and to the development of rice roots (Kartika *et al.*, 2021). However, no research has been found on how the role of biochar was given together with other organic amendments in acid sulphate soil. Therefore, it is necessary to study how the effect of biochar is given together with organic amendments to *C. odorata* and rice straw. This research is expected to contribute to improving the nature of acid sulphate soil in West Kalimantan, which has been one of the types of soil widely used as an agricultural activity for rice crop. In connection with this background and literature review above, this study aimed at knowing the role of rice husk biochar combined with amendments to *C. odorata*, rice straw and rice husk ash in improving the properties of acid sulphate soil.

MATERIALS AND METHODS

Study Area

This research was conducted from September to November 2021. The research site was carried out at the Laboratory of the Faculty of Agriculture, Universitas Panca Bhakti, Pontianak, West Kalimantan Province, Indonesia. The height of the study site is one m from above sea level, with an average temperature and humidity of 27.6°C and 82.8%, respectively. The research position is located at a latitude of 2°05' N-3° 05'S and longitude of 108°30' E-144°10' W.

Experimental Design

The study was conducted in a greenhouse and arranged in a complete randomized design (CRD) with three replications. Biochar amendments consisted of nine kinds, namely, without amendments (control), *C. odorata* (10 t/ha), rice straw (10 t/ha), rice husk biochar (10 t/ha), rice husk ash (10 t/ha), *C. odorata* (10 t/ha)+rice husk biochar (5 t/ha), rice straw (10 t/ha)+rice husk biochar (5 t/ha), *C. odorata* (10 t/ha)+rice husk ash (5 t/ha), and rice straw (10 t/ha)+rice husk ash (5 t/ha). In all, 27 polybags were needed.

Research Procedures

The material in this study used acid sulphate soil taken on the experimental land of the West Kalimantan, Sungai Kakap, Agricultural Technology Assessment Center, and land farmers' work on rice crops. This area included the development of swampland agriculture which had built irrigation canals since 1980 and was currently designated as an Integrated Farmer Business Area. Soil picking was carried out randomly at a soil depth of 0-20 cm, then mixed, air-dried, and cleaned. Then put in polybags was as much as 8 kg.

Biochar was made by taking rice husk from grain harvested from rice plants grown directly in the acid sulphate soil of the Kakap River, West Kalimantan, Indonesia. First, the rice mill was taken the grain to obtain rice husk. Furthermore, the rice husk was placed on the pyrolysis device to be processed into biochar by incomplete combustion (limited air) in a reactor made of Pertamina drums.

Organic amendments to *C. odorata* were taken from around the field, then chopped to a size of 1-2 cm. Rice straw was taken from waste harvested rice grown and grown by farmers in acid sulphate soil and chopped within 1-2 cm. Rice husks were taken from waste from rice milling, then used as ash through burning in open areas to produce grayish-white ash.

Prepared acid sulphate soil introduced into 27 polybags until it reached 8 kg. Further, the dose of soil amendment was calculated based on the soil weight in the polybag area, mixed with the soil in each polybag to a soil depth of 20 cm, after which it was incubated with moisture content until it was closed to the field capacity for 30 days.

At 30 days after incubation, from each study unit, soil sampling of about 50 g was undisturbed to determine the weight of the soil moisture content accident. In addition, a disturbed soil sample of 100 g was taken from each research unit for analysis of chemical properties.

Observation Parameters

The soil bulk density (BD) was determined by the clod method described by Blake & Harke. The total pore was calculated from the soil moisture content (v/v) at the matrix potential of 0 kPa. The available groundwater was estimated by reducing the soil moisture content in the matrix potential by -33 kPa (field capacity) with the soil moisture content in the matrix potential, which was -15 Mpa (wilting point). The soil moisture content at this potential matrix was determined using a pressure plate device. The strength of the soil was measured with a hand penetrometer (Daikie) at a soil depth of 15 cm.

The pH level was measured in a soil solution at a ratio of 1 : 2.5 (with deionized water), using a pH meter (Jenway 3305). Walkley and Black's wet oxidation method was used to determine organic C levels (Soil Survey Laboratory Staff). Total N levels were measured by the Kjeldhal (Bremner and Mulvaney) method. Al₃+and Fe₂+were extracted with 1 M

KCl (Barnhisel and Bertsch). CEC was extracted with 1 M NH₄Oac (buffer at pH 7.0), and the concentration of alkaline cations was measured using AAS (Shimadzu), P-available with Bray I.

Statistical Analysis

Statistical analysis was carried out on each observation parameter data to determine the effect of the amendments on the nature of acid sulphate soil. The data obtained were analyzed by analysis of variance (ANOVA) at 5% significant level. In addition, the test of least significant difference (LSD) at 5% significant level was used to determine the average value between treatments.

RESULTS AND DISCUSSION

Effect of Various Soil Amendments on Some Physical Soil Properties

Applying biochar improved the soil's physical properties (Aslam *et al.*, 2014). Applying biochar and organic amendments influenced the changes in some of the observed soil physical properties. The statistical test is shown in Table 1. The soil BD was lower on all amendments granted. For example, in soils fed a single rice husk biochar (10 t/ha), soil BD decreased from 1.27 mg/m³ in the control soil to 1.15 mg/cm^3 . However, not significantly different from soils was given straw (10 t/ha) and a combination of straw+biochar, *C. odorata*+rice husk ash, and straw+rice husk ash. Furthermore, in the combination of *C. odorata* (10 t/ha)+rice husk biochar (5 t/ha), a BD of 1.13 mg/m^3 was obtained and was the lowest soil BD. Still, it was not significantly different from the soil, given a combination of straw (10 t/ha) + rice husk biochar (5 t/ha) produced soil BD of 1.14 mg/m^3 . These results indicated an improvement in the physical soil properties if biochar was given together with organic amendments to *C. odorata*.

Applying various biochar amendments and their combinations increased the soil's total pore. The statistical test results are shown in Table 1. There was a significant difference in the total pore of the soil between the various amendments. Control soils had the lowest total pore (44.43%), contrasting with soils that provided various amendments. On soils with rice husk biochar (10 t/ha), the total soil pore increased to 54.21%, significantly different from other single amendments. Furthermore, a combination of *C. odorata* (10 t/ha)+rice husk biochar (5 t/ha) caused the total pore to increase to 56.73% and was the highest total pore.

The application of various amendments, both single and combined, affected the decrease in soil strength as measured as soil penetration resistance (N/cm²) at conditions of pF 0, pF 2 and pF 2.5, where the higher the soil matric suction (pF), the higher the soil penetration resistance. The under saturated conditions (pF 0), the various soil amendments did not offer a noticeable difference. In pF 2 showed that the application of rice husk biochar (10 t/ha) could reduce soil penetration resistance from 310.00 N/cm² (control soil) to 223.33 N/cm², but not significant difference between *C. odorata* (10 t/ha)+rice straw or a combination of straw (10 t/ha)+rice husk biochar (5 t/ha). Furthermore, soils

given a combination of *Cromolaena odorata* (10 t/ha)+biochar (5 t/ha) obtained the lowest penetration resistance (220.00 N/cm²), and this treatment was not significantly different from soils have given *C. odorata* (10 t/ha), rice husk biochar (10 t/ha), and a combination of rice straw (10 t/ha)+rice husk biochar (5 t/ha).

Similarly, in pF 2.5, the application of rice husk biochar (10 t/ha) could reduce the value of soil penetration resistance from 500.00 N/cm² (control soil) to 390.00 N/cm², and this treatment not significantly different from soils had given *C. odorata* (10 t/ha), rice straw (10 t/ha), rice husk ash (10 t/ha), and a combination of rice straw (10 t/ha)+rice husk ash (5 t/ha). Furthermore, the combination of *C. odorata* (10 t/ha)+rice husk biochar (5 t/ha) achieved the lowest penetration resistance (340.00 N/cm²). Also, it was not significantly different from the combination of rice straw (10 t/ha)+biochar (5 t/ha) (360.00 N/cm²). It indicated that rice husk biochar had the potential to control soil strength. In line with Das *et al.* (2021), applying biochar into the soil could reduce soil strength.

Effect of Rice Husk Biochar on Some Soil Chemical Properties

The application of several soil amendments and their combinations could influence some chemical properties of acid sulphate soil. The results of statistical tests in Table 2 show that soil reaction (pH) of the soil increased in all treatments given compared to controls. For example, in the application of rice husk biochar showed an increase in pH from 3.36 (control soil) to 4.40 on soils given biochar (10 t/ha) but did not significantly differ from other single amendments. Furthermore, the combined application of *C. odorata* (10 t/ha)+biochar (5 t/ha) obtained the highest increase in soil pH (4.58), in marked contrast to soils given rice husk ash but not significantly different from single amendments or other combinations.

Applying single and combined soil amendments affected the increase in soil C-organics. The statistical test results in Table 2 show that the control soil had the lowest C-organic (1.94%), in stark contrast to other amended soils, except that the soil had given rice husk ash (10 t/ha) was not significantly different. The soil given a single amendment of rice straw (10 t/ha) obtained 4.58% C-organic, in stark contrast to the control soil and the soil given rice husk ash, but not significantly different from other soil amendments. Similarly, soils given a combination of *C. odorata* (10 t/ha)+biochar husk (5 t/ha) achieved the highest soil C-organic (4.93%) in marked contrast to control soils and soils fed with rice husk ash but did not significantly differ from other soil amendments.

Applying various amendments and combinations to acid sulphate soil generally affected the increase in P-available. The results of statistical tests (Table 2) show that control soils had the lowest P-available content (0.21%), significantly different from soils given various amendments. In soils fed with a rice husk biochar (10 t/ha), the P-available increased to 0.32%. Still, it was not significantly different from other combined or single-amended soils, except with soils that were given rice husk

ash with a lower P-available (0.27%). The highest P-available was obtained on soils given a combination of *C. odorata* (10 t/ha)+biochar (5 t/ha) (0.34\%), in stark contrast to control soils and those topped with rice husk ash but not significantly different from single amendments or other combinations.

The decrease in Al-dd and Fe-dissolve in acid sulphate soil also occured due to applying various soil amendments and their combinations. Statistical tests (Table 2) showed that the highest Al-dd was obtained on control soil (3.84%), further decreasing and differing markedly from soils given various single and combined amendments. On single-amendment soils of rice husk biochar (10 t/ha), Al-dd decreased to 3.16%, significantly different from other single-amended soils. Furthermore, the soil given the combination of *C. odorata* (10 t/ha)+rice husk biochar (5 t/ha) obtained the lowest Al-dd content (3.02%). It was not significantly different from the soil given a combination of rice straw (10 t/ha)+rice husk biochar (5 t/ha), but it was significantly different from other single amended soils. In soluble Fe content, the application of rice husk biochar (10 t/ha) could reduce soluble Fe from 3.61% on control soil to 3.10% and was significantly different from other single amendments. And when the soil was given a combination of rice straw (10 t/ha)+rice husk biochar (5 t/ha) caused the lowest decrease in Al-dd (2.91%). Still, it was not significantly different from soil that was given a combination of *C. odorata* (10 t/ha)+rice husk biochar (5 t/ha).

The CEC of land has also increased due to the application of various land amendments and their combinations. The statistical test shown in Table 2 shows that the control soil had the lowest CEC (6.64 Cmol/kg), not significantly different from soils fed with *C. odorata* and rice straw. On the other hand, on soils provided with rice husk biochar (10 t/ha), the soil CEC increased to 8.03 Cmol/kg, but it was not significantly different from soil given rice straw and rice husk ash.

In line with the results that have been put forward, Yamato *et al.* (2006) found that the application of biochar from *Acacia mangium* bark into the soil in Sumatra led to changes in soil chemical properties through increased pH, N-total and P-available, CEC and lowered interchangeable Al. Liang *et al.* (2006) found that two mechanisms could create a larger CEC. First, a higher charged density per unit of surface acreage means a higher degree of oxidation of soil organic matter. Second, the presence of a higher surface area for cation absorption or the combined effect of both. Glaser *et al.* (2002) posit that oxidation of aromatic C and carboxyl formation were the main reasons for the high CEC. Liang *et al.* (2006) also found that the CEC per unit C was higher, and the load density was higher in carbon-black-rich Anthrosol compared to soils that were poor in carbon black. In addition, Anthrosol showed a higher surface acreage due to its higher carbon black concentration.

From the results, the great potential of rice husk biochar against improving soil properties increased when given together with the biomass *C. odorata*. It could happen because *C. odorata* contained higher humate and fulvic organic acids, so that it could encourage the complex mechanisms

of organo-mineral organs and soil aggregation. In addition, *C. odorata* can highly suppress Al and Fe, thereby increasing the P availability in the soil due to the presence of organic acids released during decomposition. Research results by Suri and Yudono (2020) also found that using *C. odorata* compost could improve soil quality and nutrient uptake of lettuce plants.

C. odorata can contribute to the care of soil C-organic and physical properties due to a reasonably high polyphenol content that could inhibit the decomposition and mineralization of N organic matter, even though it had a low C/N ratio.

CONCLUSION

In conclusion, the amendment of rice husk biochar combined with amendments to *C. odorata* and rice straw improved soil physical properties, namely, total soil pores, reducing soil content weight and strength. Its impact on soil chemical properties was that it could increase pH, C-organic, P-available, CEC, lower Al-dd and Fe soluble. Therefore, applying biochar and organic amendments to *C. odorata* and rice straw could improve the properties of acid sulphate soil. Consequently, we recommend increasing the carrying capacity of acid sulphate soil against plant growth. It can be done by enhancing soil properties through the organic amendments application of *C. odorata* or straw enriched with rice husk biochar.

ACKNOWLEDGEMENTS

The authors thank the Institute for Research and Community Service, Universitas Panca Bhakti, which has given support for research funds. In addition, they thank the Faculty of Agriculture, Universitas Panca Bhakti, which has provided loans for facilities in the form of equipment for research.

REFERENCES

- Agustina, R., Jumadi, R., Firmani, U. and Abdul Faisal, R. H. (2019). Effects of decomposition rate of *Chromolaena odorata* and straw rice in fresh and compost form to the growth and yield of rice. *IOP Conf. Ser. Earth Environ. Sci.* 250 : 1-5.
- Aslam, Z., Khalid, M. and Aon, M. (2014). Impact of biochar on soil physical properties. *Sch. J. Agric. Sci.* **4** : 280-84.
- Beusch, C. (2021). Biochar as a soil ameliorant : How biochar properties benefit soil fertility–A review. J. Geosci. Environ. Prot. 9 : 28-46.
- Bista, P., Ghimire, R., Machado, S. and Pritchett, L. (2019). Biochar effects on soil properties and wheat biomass vary with fertility management. *Agronomy* **9** : 1-10.
- Das, S. K. and Das, S. K. (2015). Acid sulphate soil : Management strategy for soil health and productivity. *Pop. Kheti* **3** : 3-8.

- Das, S., Mohanty, S., Sahu, G., Rana, M. and Pilli, K. (2021). Biochar : A sustainable approach for improving soil health and environment. In : Soil Erosion–Current Challenges and Future Perspectives in a Changing World. pp. 1-18.
- Ebido, N. E., Edeh, I. G., Unagwu, B. O., Nnadi, A. L., Ozongwu, O. V., Obalum, S. E. and Igwe, C.
 A. (2021). Rice-husk biochar effects on organic carbon, aggregate stability and nitrogen fertility of coarse-textured Ultisols evaluated using *Celosia argentea* growth. *Sains. Tanah.* 18 : 177-87.
- Gewaily, E. (2019). Impact of compost rice straw and rice straw as organic fertilizer with potassium treatments on yield and some grain quality of Giza 179 rice variety. *J. Plant Prod.* **10** : 143-51.
- Glaser, B., Lehmann, J. and Zech, W. (2002). Ameliorating physical and chemical properties of highly weathered soils in the tropics with charcoal–A review. *Biol. Fertil. Soils* **35** : 219-30.
- Jaswal, A., Mehta, C. M. and Singh, A. (2022). Probing the impact of biochar combined with organic and inorganic amendments on soil carbon pools of rice (*Oryza sativa*)-wheat (*Triticum aestivum*) cropping system. *Res. Crop* 23 : 508-15.
- Kalu, S., Simojoki, A., Karhu, K. and Tammeorg, P. (2021). Long-term effects of softwood biochar on soil physical properties, greenhouse gas emissions and crop nutrient uptake in two contrasting boreal soils. *Agric. Ecosyst. Environ.* **316** : doi.org/10.1016/j.agee.2021.107454.
- Kartika, K., Sakagami, J. I., Lakitan, B., Yabuta, S., Akagi, I., Widuri, L. I., Siaga, E., Iwanaga, H. and Nurrahma, A. H. I. (2021). Rice husk biochar effects on improving soil properties and root development in rice (*Oryza glaberrima* Steud.) exposed to drought stress during early reproductive stage. *AIMS Agric. Food* 6 : 737-51.
- Kocsis, T., Ringer, M. and Biró, B. (2022). Characteristics and applications of biochar in soil-plant systems : A short review of benefits and potential drawbacks. *Appl. Sci.* **12** : 1-16.
- Lestari, M. F. and Rachmawati, D. (2020). The effect of rice husk ash on growth of rice (*Oryza sativa* (L.) 'Sembada Merah') on salinity stress. *AIP Conf. Proc.* **2260** : 1-6.
- Liang, B., Lehmann, J., Solomon, D., Kinyangi, J., Grossman, J., O'Neill, B., Skjemstad, J. O., Thies, J., Luizão, F. J., Petersen, J. and Neves, E. G. (2006). Black carbon increases cation exchange capacity in soils. *Soil Sci. Soc. Am. J.* **70** : 1719-30.
- Mahmoud, E., Ibrahim, M., Robin, P., Akkal-Corfini, N. and El-Saka, M. (2009). Rice straw composting and its effect on soil properties. *Compost Sci. Util.* **17** : 146-50.
- Murtaza, G., Ahmed, Z., Usman, M., Tariq, W., Ullah, Z., Shareef, M., Iqbal, H., Waqas, M., Tariq, A., Wu, Y., Zhang, Z. and Ditta, A. (2021). Biochar induced modifications in soil properties and its impacts on crop growth and production. *J. Plant Nutr.* 44 : 1677-91.
- Ogunleye, K. S., Okeke, O. S., Umego, M. O. and Ogbonnaya, O. U. (2019). Extraction behaviour of cadmium in biochar amended soils. *Crop Res.* **54** : 59-64.
- Shamshuddin, J., Elisa Azura, A., Shazana, M. A. R. S., Fauziah, C. I., Panhwar, Q. A. and Naher,

U. A. (2014). Properties and management of acid sulfate soils in South-east Asia for sustainable cultivation of rice, oil palm and cocoa. *Adv. Agron.* **124** : 91-142.

- Sovova, S., Enev, V., Smilek, J., Kubikova, L., Trudicova, M., Hajzler, J. and Kalina, M. (2021). The effect of biochar application on soil properties and growth of the model plant *Zea mays*. *Ecocycles* **7** : 46-54.
- Suri, A. M. and Yudono, P. (2020). Effects of *Chromolaena odorata* compost on soil and nutrient uptake of lettuce (*Lactuca sativa*). *Planta Trop. J. Agro. Sci.* 8 : 33-38.
- Yadav, N. K., Kumar, V., Sharma, K. R., Choudhary, R. S., Butter, T. S., Singh, G., Kumar, M. and Kumar, R. (2018). Biochar and their impacts on soil properties and crop productivity : A review. *J. Pharmacogn. Phytochem.* 7 : 49-54.
- Yamato, M., Okimori, Y., Wibowo, I. F., Anshori, S. and Ogawa, M. (2006). Effects of the application of charred bark of *Acacia mangium* on the yield of maize, cowpea and peanut, and soil chemical properties in South Sumatra, Indonesia. *Soil Sci. Plant Nutr.* 52 : 489-95.
- Yin, M., Li, X., Liu, Q. and Tang, F. (2022). Rice husk ash addition to acid red soil improves the soil property and cotton seedling growth. *Sci. Rep.* **12** : 1-9.
- Yin, X., Huang, M. and Zou, Y. (2018). Potential use of biochar in raising machine-transplanted rice seedlings. *Farm. Manage.* 3 : 87-92.
- Zhang, Y., Chen, H., Xiang, J., Xiong, J., Wang, Y., Wang, Z. and Zhang, Y. (2022). Effect of ricestraw biochar application on the acquisition of rhizosphere phosphorus in acidified paddy soil. *Agronomy* 12 : 1-13.

Table 1. Effect of rice husk biochar with soil amendments on some physical properties of acid sulphate soil at one month after incubation.

BD	Total	Penetr	ation resistance	$e(N/cm^2)$
(mg/m^3)	pores (%)	pF 0	pF 2	pF 2.5
1.24 e	44.43 a	36.67 b	310.00 e	500.00 d
1.16 c	52.17 e	20.00 a	230.00 abc	403.33 bc
1.17 bc	53.27 f	16.67 a	243.33 bc	393.33 b
1.15 b	54.21 g	14.33 a	223.33 ab	390.00 b
1.19 d	47.30 b	20.00 a	270.00 d	403.33 bc
1.13 a	56.73 i	10.00 a	220.00 a	340.00 a
	BD (mg/m ³) 1.24 e 1.16 c 1.17 bc 1.15 b 1.19 d 1.13 a	BDTotal(mg/m³)pores (%)1.24 e44.43 a1.16 c52.17 e1.17 bc53.27 f1.15 b54.21 g1.19 d47.30 b1.13 a56.73 i	BDTotalPenetr(mg/m³)pores (%)pF 01.24 e44.43 a36.67 b1.16 c52.17 e20.00 a1.17 bc53.27 f16.67 a1.15 b54.21 g14.33 a1.19 d47.30 b20.00 a1.13 a56.73 i10.00 a	BD Total Penetration resistance (mg/m ³) pores (%) pF 0 pF 2 1.24 e 44.43 a 36.67 b 310.00 e 1.16 c 52.17 e 20.00 a 230.00 abc 1.17 bc 53.27 f 16.67 a 243.33 bc 1.15 b 54.21 g 14.33 a 223.33 ab 1.19 d 47.30 b 20.00 a 270.00 d 1.13 a 56.73 i 10.00 a 220.00 a

Rice straw (10 t/ha) + rice	1.14 ab	55.57 h	13.33 a	223.00 ab	360.00 a
husk biochar (5 t/ha)					
C. odorata (10 t/ha) + rice	1.15 b	48.90 c	16.67 a	240.00 bc	420.00 c
husk ash (5 t/ha)					
Rice straw (10 t/ha) + rice	1.16 bc	50.30 d	20.00 a	246.67 с	403.33 bc
husk ash (5 t/ha)					

Remarks: The average numbers followed by the same letter in the same column show no significant difference in the LSD test at 5% significant level.

Tab	le 2.	Effect	of	rice	husk	biochar	with	soil	amendments	on	some	chemical	properties	of	acid
sulp	hate	soil at o	one	mon	th aft	er incuba	ation.								

		C-	Р-	Al-dd	Soluble	CEC
Soil amendments	pH H ₂ O	organic	available	(me/100	Fe	(me/100
		(%)	(%)	g)	(%)	g)
Control	3.36 a	1.94 a	0.21 a	3.84 e	3.61 d	6.64 a
C. odorata (10 t/ha)	4.06 bc	4.22 b	0.29 bc	3.31 c	3.28 c	7.15 ab
Rice straw (10 t/ha)	4.28 bc	4.58 b	0.30 bc	3.42 cd	3.34 cd	7.32 abc
Rice husk biochar (10 t/ha)	4.40 bc	4.09 b	0.32 cd	3.16 b	3.10 abc	8.03 c
Rice husk ash (10 t/ha)	3.98 b	2.78 a	0.28 b	3.51 d	3.34 cd	7.76 bc
C. odorata $(10 \text{ t/ha}) + \text{rice}$	4.58 c	4.93 b	0.34 d	3.02 a	2.94 ab	10.04 e
husk biochar (5 t/ha)						
Rice straw (10 t/ha) + rice	4.48 bc	4.75 b	0.33 d	3.08 ab	2.91 a	9.67 de
husk biochar (5 t/ha)						
C. odorata $(10 \text{ t/ha}) + \text{rice}$	4.26 bc	4.70 b	0.31 cd	3.37 c	3.25 c	9.17 d
husk ash (5 t/ha)						
Rice straw (10 t/ha) + rice	4.26 bc	4.73 b	0.31 cd	3.47 d	3.22 bc	9.20 d
husk ash (5 t/ha)						

Remarks: The average numbers followed by the same letter in the same column show no significant difference in the LSD test at 5% significant level.

Persetujuan artikel untuk dipublikasi : 8 Desember 2022



Online publication : 16 Desember 2022

M Pentin	ig - agusalim@upb.ac.id - E	× +		· · · · · · · · · · · · · · · · · · ·	/ - 0
$\leftarrow \rightarrow$	C 🗎 mail.google.c	com/mail/u/0/?t	ab=rm&ogbl#imp		ê 🖈 🗖 🍥
YouTub	e 🛃 Maps				
=	M Gmail		Q is:important	× 幸 (• Aktif • ⑦ ③ Ⅲ	🎯UPB 🎯
Mail	🖉 Tulis		Dari 👻 Kapan saja 👻	Ada lampiran Kepada 👻 Belum dibaca Penelusuran lanjutan	
	Kotak Masuk	5.342	□ • C :	1–50 dari 96	< >
Chat	📥 Berbintang		🔲 🚖 Gaurav, saya 2	Kotak Masuk Online Publication of manuscript ROC-907 " Role of biochar amendments in improving the properties o	16 Des
:0:	③ Ditunda		🗌 🚖 VEDPAL, saya 3	Kotak Masuk Galley copy of paper for publication (10. 11147) - Thanks for your confirmation to consider galley copy f	8 Des
Spaces	Terkirim			10. 11147 Galley	
	🗋 Draf	4	🔲 🕁 Jurnal Ilmu Pertani.	Kotak Masuk [JIPI] New notification from Jurnal Ilmu Pertanian Indonesia - You have a new notification from Jurnal Il	7 Des
Weet	~ Ciutkan		🗆 🕁 Jurnal Ilmu Pertani.	Kotak Masuk [JIP] New notification from Jurnal Ilmu Pertanian Indonesia - You have a new notification from Jurnal II	6 Des
	Penting			Tract track f UNI Descurred Peacet. Your accounced has been auroacted fully react for use with the Scientific Journals	E Daa
	I Semua Email			Kotak Masuk [JIP] Password Reset - Your password has been successfully reset for use with the scientific Journals	5 Des
	Spam		📋 ☆ Dr. Iman Rusmana 2	Kotak Masuk [JIPI] Password Reset Confirmation - We have received a request to reset your password for the Scienti	5 Des
	i Sampah		🗄 🗖 🛧 Scholastica	Kotak Masuk [The Abstract] Author support insights + new publishing analytics - In this issue: Authorship 🗈 👔	⊠ ()
	► D Kategori		🔲 🚖 Editorial, saya 2	Kotak Masuk Successful Completion from Initial Scrutiny BMS-TOASJ-2022-63 Forwarded message	18 Nov
	🔅 Kelola label		🗌 🚖 Billing Department	Kotak Masuk Figure Improvement Service BMS-TOASJ-2022-63 - Ref #: 115922 Dear Dr, Agusalim Masulili This is wit	14 Nov
	+ Buat label bar	u	🔲 🕁 Bentham, saya 3	Kotak Masuk Figure Improvement Query BMS-TOASJ-2022-63 - PROOF OF PAPER To Open Agriculture Journal Dea	14 Nov
	Label	+		🔯 Manuscript_Ag) 🔤 115922_lab_rep	
			🔲 🚖 saya, Jurnal 7	Kotak Masuk Mohon Informasi - Yth. Editor Mohon perkenan informasi, kelanjutan review artikel kami. Terima kasih b	14 Nov



https://gauravpublications.com/journal/research-on-crops/ROC-907



Final copy of final published paper (ID 11147) :19 Desember 2022.

\rightarrow	C I	mail.google.com	n/mail/u/0/?ta	ab=rm&ogbl#imp	کا 🚸	👌 🏚 🖬 🎯
YouTub	e 🛃	Maps				
=	M	Gmail		Q is:important	× 荘 ● Aktif ▼ ⑦ 參 ⅲ	🎯 UPB 🤡
Mail	0	Tulis		Dari 👻 Kapan saja 👻	Ada lampiran Kepada 💌 Belum dibaca Penelusuran lanjutan	
		Kotak Masuk	5.342	□ • C :	1–50 dari 96	< >
Chat	☆	Berbintang		🗆 🚖 VEDPAL, saya 2	Kotak Masuk pdf copy of final published paper (ID 11147) - Dear, Vedpal Singh Editor - in-Chief Thank you for the info	19 Des
8	0	Ditunda			🚾 11. 787-794.pdf	
aces		Terkirim		🔲 🚖 Gaurav, saya 2	Kotak Masuk Online Publication of manuscript ROC-907 " Role of biochar amendments in improving the properties o	16 Des
⊡• leet		Draf	4	VEDPAL, sava 3	Kotak Masuk Galley copy of paper for publication (10, 11147) - Thanks for your confirmation to consider galley copy f	8 Des
	ŝ	Penting			10. 11147 Galley	
	Fe	Terjadwal		D	Participation I UPD New antification from Averal line Partacian Independent Ver have a new antification from Averal II	7.000
	e	Semua Email			Katak Masuk [JiPi] Yew Notification from Juman limb Pertaman indonesia - You have a new notification from Juman li	7 Dens
	٢	Spam		🔄 🚖 Jurnal Ilmu Pertani.	Kotak Masuk [JIPI] New notification from Jurnal Ilmu Pertanian Indonesia - You have a new notification from Jurnal II	6 Des
	Ē	Sampah		🔲 🚖 Dr. Iman Rusmana	Kotak Masuk [JIPI] Password Reset - Your password has been successfully reset for use with the Scientific Journals	5 Des
	• 🗅	Kategori		📋 🚖 Dr. Iman Rusmana 2	Kotak Masuk [JIPI] Password Reset Confirmation - We have received a request to reset your password for the Scienti	5 Des
	\$	Kelola label		🗌 🚖 Scholastica	Kotak Masuk [The Abstract] Author support insights + new publishing analytics - In this issue: Authorship dos and do	30 Nov
	+	Buat label baru		🗌 🚖 Editorial, saya 2	Kotak Masuk Successful Completion from Initial Scrutiny BMS-TOASJ-2022-63 Forwarded message	18 Nov
	Lab	el	+	🔲 🚖 Billing Department	Kotak Masuk Figure Improvement Service BMS-TOASJ-2022-63 - Ref #: 115922 Dear Dr, Agusalim Masulili This is wit	14 Nov
				🔲 🕁 Bentham, saya 3	Kotak Masuk Figure Improvement Query BMS-TOASJ-2022-63 - PROOF OF PAPER To Open Agriculture Journal Dea	14 Nov



Complete Publication of manuscript no. ROC-907 entitled Role of biochar amendments in improving the properties of acid sulphate soil.

Publication : 25 Desember 2022.

https://gauravpublications.com/journal/research-on-crops/ROC-907

	C A mail accede com/	G effect h	umat acid on soil - Googi 🗙 G sci hub -	Google Search X M Penting - agusalim@upb.ac.id - X +	
7 1	C Main.google.com/r	naii/u/3/:0	goiminip		© # L 🐡
	M Gmail		Q is:important	× ≇ ● Aktif ▼ ⑦ 戀 Ⅲ	🛞 UPB 🌀
9 •	0 Tulis		Dari 🔹 Kapan saja 👻	Ada lampiran Kepada 💌 Belum dibaca Penelusuran lanjutan	
	😡 Kotak Masuk	5.343	□ • ♂ :	1–50 dari 97	$\langle \rangle$
it 1	☆ Berbintang③ Ditunda		🔲 🚖 Gaurav Publications	Kotak Masak Complete Publication of manuscript no. ROC-907 entitled Role of biochar amendments in improving the ROC.jpg III. 787-794.pdf III Contents ROC V	22.25
es	► Terkirim		🔲 👷 Jurnal Ilmu Pertani.	Kotak Masuk [JIPI] New notification from Jurnal Ilmu Pertanian Indonesia - You have a new notification from Jurnal Ilm	22 Des
i St	Ciutkan	4	🗌 🚖 VEDPAL, saya 2	Kotak Masuk pdf copy of final published paper (ID 11147) - Dear, Vedpal Singh Editor - in-Chief Thank you for the infor	19 Des
	Penting				
	Fo Terjadwal		🔲 🛧 Gaurav, saya 2	Kotak Masuk Online Publication of manuscript ROC-907 " Role of biochar amendments in improving the properties of	16 Des
	Semua Email		🔲 🚖 VEDPAL, saya 3	Kotak Masuk Galley copy of paper for publication (10. 11147) - Thanks for your confirmation to consider galley copy fo	8 Des
	III Sampah ▶ D Kategori		📋 🚖 Jurnal Ilmu Pertani.	Kotak Masuk [JIPI] New notification from Jurnal Ilmu Pertanian Indonesia - You have a new notification from Jurnal Ilm	7 Des
	Kelola label		🔲 🚖 Jurnal Ilmu Pertani.	Kotak Masuk [JIPI] New notification from Jurnal Ilmu Pertanian Indonesia - You have a new notification from Jurnal Ilm	6 Des
	+ Buat label baru		🔲 🚖 Dr. Iman Rusmana	Kotak Masuk [JIPI] Password Reset - Your password has been successfully reset for use with the Scientific Journals o	5 Des
	Label	+	🔲 🚖 Dr. Iman Rusmana 2	Kotak Masuk [JIPI] Password Reset Confirmation - We have received a request to reset your password for the Scienti	5 Des
			🖸 🕁 Scholastica	Kotak Masuk [The Abstract] Author support insights + new publishing analytics - In this issue: Authorship dos and do	30 Nov
			🗄 🗖 🛧 Editorial, saya 2	Kotak Masuk, Successful Completion from Initial Scrutiny BMS-TOASJ-2022-63 Forwarded me 🔳 👔	0



Final Publish in web:

https://gauravpublications.com/journal/research-on-crops/volume-23/issue-4december

Gaurav Publications	Journals	Services 🗸	Policies 🗸	📥 LOGIN
Home	Research On Crops > Journal > Research on Crops > VOLUME 23 > ISSUE	4 (DECEMBER)		
VOLUME 23, ISSUE 4 (DECE	EMBER)			Print 6019 5070-2006 Octow 5388 2540-74-2
ISSN: 2348-7542 (Online) 0972-3226 (Print) I.F. : IF 2012: 0.103 Naas Rating: 4.56 SJR: 0.27			Research of	n Grops
In this issue (29 articles)				
Exogenous melatonin application mitigates anthesis stages in rice (Oryza sativa) R. MEGALA, M. K. KALARANI, P. JEYAKUMAR, N. SEI	the adverse impact of drought stress at panicle i	nitiation and		
DOI :10.31830/2348-7542.2022.ROC-836 Article Id :RC	DC-836	Pages 711-718		
Supplementary potassium application enric A. K. M. GOLAM SARWAR, MD. SABIBUL HAQUE, MD	ched micronutrient status of purple rice (Oryza se . ZAKIR HOSSEN AND A. K. M. AHSAN KABIR	ativa L.)	Agricultural Research Inf Game Fellowing Hisar, Ind	ormation Centre mati
DOI :10.31830/2348-7542.2022.ROC-871 Article Id :RC	DC-871	Pages 719-728		
Modification of root architecture of rice by stress	guard cell and rhizosphere halotolerant bacteria	under saline	→ Editorial Board → Aims and Scope	
DOI :10.31830/2348-7542.2022.ROC-835 Article Id :RC	IC-835	Pages 729-736	→ Role of Reviewers	
			→ Editorial Policy	

Gaurav Publications Journals Services 🗸	Policies 🗸		
Influence of basic tillage and coulisse strips on the weed infestation and yield of spring wheat aestivum) in the Tyumen region N. FISUNOV, O. SHULEPOVA AND N. SANNIKOVA	(Triticum	Search Article	
DOI :10.31830/2348-7542.2022.ROC-905 Article Id :ROC-905	Pages 770-775		
Pffeat of all at density and falling and of sing an exact time month of any (7.5 month)		Contact Us	
AIYMEN AHMED A. K. AL-ABASSI, WAEL F. A. AL-SHAMARY, KHAMAEL ALI KAREEM AND ABDEL MONNEM SADALA	AHA KAHLEI	Enter Name	
DOI :10.31830/2348-7542.2022.ROC-827 Article Id :ROC-827	Pages 776-780	Enter Email	
Effect of addition of biopolymer coated DAP on phosphorus release kinetics over an incubatio M DHANALAKSHMI S MEENA P JANAKI S KARTHIKEYAN AND L NALINA	on time	Select Country 🗸	
DOI :10.31830/2348-7542.2022.ROC-906 Article Id :ROC-906	Pages 781-786	Message	
Role of biochar amendments in improving the properties of acid sulphate soil Agusalim Masulili, Sutikarini, Rini Suryani, Ida ayu Suci, Ismail Astar, Hardi dominikus Bancin and Dal to appropriate Zero 2000 per set. 1 Attick 14:000-007	PAIMAN	4	
Doi:10.31830/2348-7542.2022.ROC-907 Article in :ROC-907	Pages 101-154	(
Physiological and phyto-pathological assessment scion-rootstock combinations for apple cv. A M. sieversii	Aport and	Saya bukan robot	
SAGI SOLTANBEKOV, SVETLANA DOLGIKH, MOLDIR ZHUMAGULOVA, AIGUL MADENOVA, ZHANNA ISINA AND BAL KABYLBEKOVA	LNUR	Submit	
DOI :10.31830/2348-7542.2022.ROC-837 Article Id :ROC-837	Pages 795-800		
Residual effect of biofortified iodine in soil, plant, crop yield and quality of tomato (Solanum lycopersicum L.)			

https://gauravpublications.com/journal/research-on-crops/volume-23/issue-4-

december/ROC-907

